

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 August 2002 (29.08.2002)

PCT

(10) International Publication Number
WO 02/066481 A1

(51) International Patent Classification⁷: C07D 487/04, A61K 31/5025, A61P 35/00

(74) Agents: **BILL, Kevin** et al.; AstraZeneca, Global Intellectual Property, Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).

(21) International Application Number: PCT/GB02/00603

(22) International Filing Date: 12 February 2002 (12.02.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0103926.2 17 February 2001 (17.02.2001) GB

(71) Applicant (for all designated States except MG, US): **ASTRAZENECA AB** [SE/SE]; Sodertalje, S-151 85 (SE).

(71) Applicant (for MG only): **ASTRAZENECA UK LIMITED** [GB/GB]; 15 Stanhope Gate, London, Greater London W1Y 6LN (GB).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **THOMAS, Andrew, Peter** [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).

(81) Designated States (national): AE, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PYRIMIDINE DERIVATIVES FOR INHIBITION OF CELL-PROLIFERATION



(I)

(57) Abstract: Compounds of the formula (I): wherein Ring A, R¹, R³, R⁴, p, q, and n are as defined within and a pharmaceutically acceptable salts and *in vivo* hydrolysable esters are described. Also described are processes for their preparation and their use as medicaments, particularly medicaments for producing a cell cycle inhibitory (anti-cell-proliferation) effect in a warm-blooded animal, such as man.

WO 02/066481 A1

PYRIMIDINE DERIVATIVES FOR INHIBITION OF CELL-PROLIFERATION

The invention relates to pyrimidine derivatives, or pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof, which possess cell-cycle inhibitory activity and are accordingly useful for their anti-cell-proliferation (such as anti-cancer) activity and are therefore useful in methods of treatment of the human or animal body. The invention also relates to processes for the manufacture of said pyrimidine derivatives, to pharmaceutical compositions containing them and to their use in the manufacture of medicaments of use in the production of an anti-cell-proliferation effect in a warm-blooded animal such as man.

A family of intracellular proteins called cyclins play a central role in the cell cycle. The synthesis and degradation of cyclins is tightly controlled such that their level of expression fluctuates during the cell cycle. Cyclins bind to cyclin-dependent serine/threonine kinases (CDKs) and this association is essential for CDK (such as CDK1, CDK2, CDK4 and/or CDK6) activity within the cell. Although the precise details of how each of these factors combine to regulate CDK activity is poorly understood, the balance between the two dictates whether or not the cell will progress through the cell cycle.


The recent convergence of oncogene and tumour suppressor gene research has identified regulation of entry into the cell cycle as a key control point of mitogenesis in tumours. Moreover, CDKs appear to be downstream of a number of oncogene signalling pathways. Disregulation of CDK activity by upregulation of cyclins and/or deletion of endogenous inhibitors appears to be an important axis between mitogenic signalling pathways and proliferation of tumour cells.

Accordingly it has been recognised that an inhibitor of cell cycle kinases, particularly inhibitors of CDK2, CDK4 and/or CDK6 (which operate at the S-phase, G1-S and G1-S phase respectively) should be of value as a selective inhibitor of cell proliferation, such as growth of mammalian cancer cells.

The present invention is based on the discovery that certain pyrimidine compounds surprisingly inhibit the effects of cell cycle kinases showing selectivity for CDK2, CDK4 and CDK6, and thus possess anti-cell-proliferation properties. Such properties are expected to be of value in the treatment of disease states associated with aberrant cell cycles and cell proliferation such as cancers (solid tumours and leukemias), fibroproliferative and differentiative disorders, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma,


acute and chronic nephropathies, atheroma, atherosclerosis, arterial restenosis, autoimmune diseases, acute and chronic inflammation, bone diseases and ocular diseases with retinal vessel proliferation.

Accordingly, the present invention provides a compound of formula (I):



wherein:

Ring A is a group of formula (IA) or (IB):



wherein one or more of X^1 , X^2 , X^3 and X^4 are nitrogen and the others are CR^5 wherein the values of R^5 may be the same or different;

R^1 is halo, nitro, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, amino, carboxy, carbamoyl, mercapto, sulphonamoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkylS(O) $_a$ wherein a is 0 to 2, N -(C_{1-6} alkyl)sulphonamoyl or N,N -(C_{1-6} alkyl) $_2$ sulphonamoyl; wherein R^1 may be optionally substituted on carbon by one or more R^6 ;

R^2 and R^5 are independently of one other selected from hydrogen, halo, nitro, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, amino, carboxy, carbamoyl, mercapto, sulphonamoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkylS(O) $_a$ wherein a is 0 to 2, C_{1-6} alkoxycarbonyl, N -(C_{1-6} alkyl)sulphonamoyl, N,N -(C_{1-6} alkyl) $_2$ sulphonamoyl, phenyl, heterocyclic group, phenylthio or (heterocyclic group)thio; wherein any R^2 or R^5 may be optionally substituted on carbon by

one or more R^7 ; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^8 ;

n is 0 to 2, wherein the values of R^1 may be the same or different;

R^3 is halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl,

5 C_{2-6} alkenyl or C_{2-6} alkynyl;

p is 0-4; wherein the values of R^3 may be the same or different;

R^4 is a group A-E-; wherein

A is selected from C_{1-6} alkyl, phenyl, a heterocyclic group, C_{3-8} cycloalkyl, phenyl C_{1-6} alkyl, (heterocyclic group) C_{1-6} alkyl or C_{3-8} cycloalkyl C_{1-6} alkyl; wherein A may be
10 optionally substituted on carbon by one or more R^9 ; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^{10} ;

E is a direct bond or -O-, -C(O)-, -OC(O)-, -C(O)O-, -N(R^a)C(O)-, -C(O)N(R^a)-, -N(R^a)-, -S(O) $_a$ -, -SO₂N(R^a)- or -N(R^a)SO₂-; wherein R^a is hydrogen or C_{1-6} alkyl optionally
15 substituted by one or more R^{11} and a is 0-2;

R^9 is independently selected from oxo, halo, nitro, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkanoylamino, N -(C_{1-6} alkyl)carbamoyl,
20 N,N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkylS(O) _{a} wherein a is 0 to 2, C_{1-6} alkoxycarbonyl, C_{1-6} alkoxycarbonylamino, benzyloxycarbonylamino, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl)₂sulphamoyl; wherein R^9 may be optionally substituted on carbon by one or more R^{12} ;

q is 0-2; wherein the values of R^4 maybe the same or different; and wherein $p + q \leq 5$;

25 R^6 , R^7 , R^{11} and R^{12} are independently selected from halo, nitro, cyano, hydroxy, trifluoromethoxy, trifluoromethyl, amino, carboxy, carbamoyl, mercapto, sulphamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxyl, methylamino, ethylamino, dimethylamino, diethylamino, N -methyl- N -ethylamino, acetylamino, N -methylcarbamoyl, N -ethylcarbamoyl, N,N -dimethylcarbamoyl, N,N -diethylcarbamoyl, N -methyl- N -ethylcarbamoyl, methylthio,
30 ethylthio, methylsulphinyl, ethylsulphinyl, mesyl, ethylsulphonyl, methoxycarbonyl, ethoxycarbonyl, N -methylsulphamoyl, N -ethylsulphamoyl, N,N -dimethylsulphamoyl, N,N -diethylsulphamoyl or N -methyl- N -ethylsulphamoyl; and

R⁸ and R¹⁰ are independently selected from C₁₋₄alkyl, C₁₋₄alkanoyl, C₁₋₄alkylsulphonyl, C₁₋₄alkoxycarbonyl, carbamoyl, N-(C₁₋₄alkyl)carbamoyl, N,N-(C₁₋₄alkyl)carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl; or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

5 In another aspect of the invention there is provided a compound of formula (I) as defined above, but wherein A is selected from C₁₋₆alkyl, phenyl, a heterocyclic group, C₃₋₈cycloalkyl, phenylC₁₋₆alkyl, (heterocyclic group)C₁₋₆alkyl or C₃₋₈cycloalkylC₁₋₆cycloalkyl; wherein A may be optionally substituted on carbon by one or more R⁹; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a
10 group selected from R¹⁰.

In this specification the term "alkyl" includes both straight and branched chain alkyl groups but references to individual alkyl groups such as "propyl" are specific for the straight chain version only. For example, "C₁₋₆alkyl" includes C₁₋₄alkyl, C₁₋₃alkyl, propyl, isopropyl and *t*-butyl. However, references to individual alkyl groups such as 'propyl' are specific for
15 the straight chained version only and references to individual branched chain alkyl groups such as 'isopropyl' are specific for the branched chain version only. A similar convention applies to other radicals, for example "phenylC₁₋₆alkyl" includes phenylC₁₋₄alkyl, benzyl, 1-phenylethyl and 2-phenylethyl. The term "halo" refers to fluoro, chloro, bromo and iodo.

Where optional substituents are chosen from "one or more" groups it is to be
20 understood that this definition includes all substituents being chosen from one of the specified groups or the substituents being chosen from two or more of the specified groups.

A "heterocyclic group" is a saturated, partially saturated or unsaturated, mono or bicyclic ring containing 4-12 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked,
25 wherein a -CH₂- group can optionally be replaced by a -C(O)- and a ring sulphur atom may be optionally oxidised to form the S-oxide(s). Examples and suitable values of the term "heterocyclic group" are morpholino, piperidyl, pyridyl, pyranyl, pyrrolyl, isothiazolyl, indolyl, quinolyl, thienyl, 1,3-benzodioxolyl, thiadiazolyl, piperazinyl, thiazolidinyl, pyrrolidinyl, thiomorpholino, pyrrolinyl, homopiperazinyl, 3,5-dioxapiperidinyl,
30 tetrahydropyranyl, imidazolyl, pyrimidyl, pyrazinyl, pyridazinyl, isoxazolyl, 4-pyridone, 1-isoquinolone, 2-pyrrolidone and 4-thiazolidone. Further examples and suitable values of the term "heterocyclic group" are morpholino, piperidyl, pyridyl, pyranyl, pyrrolyl, pyrazolyl, isothiazolyl, indolyl, quinolyl, thienyl, 1,3-benzodioxolyl, thiadiazolyl, piperazinyl,

thiazolidinyl, pyrrolidinyl, thiomorpholino, pyrrolinyl, homopiperazinyl, 3,5-dioxapiperidinyl, tetrahydropyranyl, imidazolyl, pyrimidyl, pyrazinyl, pyridazinyl, isoxazolyl, 4-pyridone, 1-isoquinolone, 2-pyrrolidone and 4-thiazolidone, particularly morpholino, pyrazolyl and pyrrolidinyl. Preferably a "heterocyclic group" is a saturated, partially saturated or
 5 unsaturated, mono or bicyclic ring containing 5 or 6 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, it may, unless otherwise specified, be carbon or nitrogen linked, a -CH₂- group can optionally be replaced by a -C(O)- and a ring sulphur atom may be optionally oxidised to form the S-oxides.

An example of "C₁₋₆alkanoyloxy" is acetoxy. Examples of "C₁₋₆alkoxycarbonyl"
 10 include methoxycarbonyl, ethoxycarbonyl, *n*- and *t*-butoxycarbonyl. Examples of "C₁₋₆alkoxy" include methoxy, ethoxy and propoxy. Examples of "C₁₋₆alkanoylamino" include formamido, acetamido and propionylamino. Examples of "C₁₋₆alkylS(O)_a wherein a is 0 to 2" include methylthio, ethylthio, methylsulphinyl, ethylsulphinyl, mesyl and ethylsulphonyl. Examples of "C₁₋₆alkanoyl" include propionyl and acetyl. Examples of
 15 "*N*-(C₁₋₆alkyl)amino" include methylamino and ethylamino. Examples of "*N,N*-(C₁₋₆alkyl)₂amino" include di-*N*-methylamino, di-(*N*-ethyl)amino and *N*-ethyl-*N*-methylamino. Examples of "C₂₋₆alkenyl" are vinyl, allyl and 1-propenyl. Examples of "C₂₋₆alkynyl" are ethynyl, 1-propynyl and 2-propynyl. Examples of "*N*-(C₁₋₆alkyl)sulphamoyl" are *N*-(methyl)sulphamoyl and *N*-(ethyl)sulphamoyl. Examples of
 20 "*N*-(C₁₋₆alkyl)₂sulphamoyl" are *N,N*-(dimethyl)sulphamoyl and *N*-(methyl)-*N*-(ethyl)sulphamoyl. Examples of "*N*-(C₁₋₆alkyl)carbamoyl" are methylaminocarbonyl and ethylaminocarbonyl. Examples of "*N,N*-(C₁₋₆alkyl)₂carbamoyl" are dimethylaminocarbonyl and methylethylaminocarbonyl. Examples of "C₃₋₈cycloalkyl" are cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. Examples of
 25 "(heterocyclic group)C₁₋₆alkyl" include pyridylmethyl, 3-morpholinopropyl and 2-pyrimid-2-ylethyl. Examples of "(heterocyclic group)thio" include pyridylthio, 3-morpholiniothio and 2-pyrimid-2-ylthio. Examples of "C₃₋₈cycloalkylC₁₋₆alkyl" are cyclopropylethyl, cyclobutylmethyl, 2-cyclopropylpropyl and cyclohexylethyl. Examples of "phenylC₁₋₆alkyl" are phenethyl, benzyl and 2-phenylpropyl.

30 A suitable pharmaceutically acceptable salt of a compound of the invention is, for example, an acid-addition salt of a compound of the invention which is sufficiently basic, for example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic, citric or maleic acid. In

addition a suitable pharmaceutically acceptable salt of a compound of the invention which is sufficiently acidic is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an organic base which affords a physiologically-acceptable cation, for example a salt with
5 methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

An *in vivo* hydrolysable ester of a compound of the formula (I) containing carboxy or hydroxy group is, for example, a pharmaceutically acceptable ester which is hydrolysed in the human or animal body to produce the parent acid or alcohol. Suitable pharmaceutically
10 acceptable esters for carboxy include C₁₋₆alkoxymethyl esters for example methoxymethyl, C₁₋₆alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl esters, C₃₋₈cycloalkoxycarbonyloxyC₁₋₆alkyl esters for example 1-cyclohexylcarbonyloxyethyl, 1,3-dioxolen-2-onylmethyl esters for example 5-methyl-1,3-dioxolen-2-onylmethyl; and C₁₋₆alkoxycarbonyloxyethyl esters for example 1-methoxycarbonyloxyethyl and may be
15 formed at any carboxy group in the compounds of this invention.

An *in vivo* hydrolysable ester of a compound of the formula (I) containing a hydroxy group includes inorganic esters such as phosphate esters and α -acyloxyalkyl ethers and related compounds which as a result of the *in vivo* hydrolysis of the ester breakdown to give the parent hydroxy group. Examples of α -acyloxyalkyl ethers include acetoxymethoxy and
20 2,2-dimethylpropionyloxy-methoxy. A selection of *in vivo* hydrolysable ester forming groups for hydroxy include alkanoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and *N*-(dialkylaminoethyl)-*N*-alkylcarbamoyl (to give carbamates), dialkylaminoacetyl and carboxyacetyl. Examples of substituents on benzoyl include morpholino and piperazino linked
25 from a ring nitrogen atom via a methylene group to the 3- or 4- position of the benzoyl ring.

Some compounds of the formula (I) may have chiral centres and/or geometric isomeric centres (E- and Z- isomers), and it is to be understood that the invention encompasses all such optical, diastereoisomers and geometric isomers that possess CDK inhibitory activity.

The invention relates to any and all tautomeric forms of the compounds of the formula
30 (I) that possess CDK inhibitory activity. In particular the skilled reader will appreciate that when R⁴ is hydrogen, the imidazole ring as drawn in formula (I) may tautomerise.

It is also to be understood that certain compounds of the formula (I) can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which possess CDK inhibitory activity.

5 Preferred values for the variable groups are as follows. Such values may be used where appropriate with any of the definitions, claims or embodiments defined hereinbefore or hereinafter.

In one aspect of the invention, preferably Ring A is a group of formula (IA).

In a further aspect of the invention, preferably Ring A is a group of formula (IB).

10 Preferably one of X^1 , X^2 , X^3 and X^4 are nitrogen and the others are CR^5 wherein the values of R^5 may be the same or different.

In one aspect of the invention, preferably X^1 is nitrogen and the others are CR^5 wherein the values of R^5 may be the same or different.

15 In another aspect of the invention, preferably X^2 is nitrogen and the others are CR^5 wherein the values of R^5 may be the same or different.

In an additional aspect of the invention, preferably X^3 is nitrogen and the others are CR^5 wherein the values of R^5 may be the same or different.

In an additional further aspect of the invention, preferably X^4 is nitrogen and the others are CR^5 wherein the values of R^5 may be the same or different.

20 Preferably R^1 is halo or $C_{1-6}alkylS(O)_a$ wherein a is 0; wherein R^1 may be optionally substituted on carbon by one or more R^6 ; wherein R^6 is hydroxy.

More preferably R^1 is bromo or 2-hydroxyethylthio.

Particularly R^1 is bromo or 2-hydroxyethylthio and n is 0-1.

Preferably R^2 is hydrogen.

25 In another aspect of the invention, preferably R^2 is selected from hydrogen, $C_{1-6}alkyl$ or $N,N-(C_{1-6}alkyl)_2amino$.

In another aspect of the invention, more preferably R^2 is selected from hydrogen, methyl, ethyl or N,N -dimethylamino.

Preferably R^5 is hydrogen.

30 In another aspect of the invention, preferably R^5 is selected from hydrogen, $C_{1-6}alkyl$ or $C_{1-6}alkoxy$; wherein R^5 may be optionally substituted on carbon by one or more R^7 ; wherein R^7 is selected from halo.

In another aspect of the invention, more preferably R^5 is selected from hydrogen, methyl, methoxy or 2,2,2-trifluoroethoxy.

In one aspect of the invention, preferably n is 2, wherein the values of R^1 may be the same or different.

5 In a further aspect of the invention, preferably n is 1.

In an additional aspect of the invention, preferably n is 0.

Preferably R^3 is sulphamoyl.

In another aspect of the invention, preferably R^3 is sulphamoyl or halo.

In another aspect of the invention, more preferably R^3 is sulphamoyl or fluoro.

10 Preferably p is 0 or 1.

Preferably R^4 is a group A-E-; wherein

A is selected from C_{1-6} alkyl; wherein A may be optionally substituted on carbon by one or more R^9 ;

15 E is $-N(R^a)C(O)-$, $-S(O)_a-$, $-SO_2N(R^a)-$ or $-N(R^a)SO_2-$; wherein R^a is hydrogen or C_{1-6} alkyl and a is 0-2;

R^9 is independently selected C_{1-6} alkoxy and N,N -(C_{1-6} alkyl)₂amino.

More preferably R^4 is a group A-E-; wherein

A is selected from methyl, ethyl or propyl; wherein A may be optionally substituted on carbon by one or more R^9 ;

20 E is $-S(O)_a-$ or $-NH SO_2-$;

R^9 is independently selected from methoxy and N,N -dimethylamino.

Particularly R^4 is N -(2-dimethylaminoethyl)sulphamoyl,

N -(2-methoxyethyl)sulphamoyl, N -methylsulphamoyl or

N -(3-dimethylaminopropyl)sulphamoyl.

25 In another aspect of the invention, preferably R^4 is a group A-E-; wherein

A is selected from C_{1-6} alkyl, C_{3-8} cycloalkyl, (heterocyclic group) C_{1-6} alkyl or C_{3-8} cycloalkyl C_{1-6} alkyl; wherein A may be optionally substituted on carbon by one or more R^9 ;

E is $-N(R^a)SO_2-$; wherein R^a is hydrogen or C_{1-6} alkyl;

30 R^9 is independently selected from hydroxy, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, N,N -(C_{1-6} alkyl)₂amino or C_{1-6} alkyl $S(O)_a$ wherein a is 0.

In another aspect of the invention, more preferably R^4 is a group A-E-; wherein

A is selected from methyl, ethyl, propyl, isopropyl, *t*-butyl, cyclopropyl, cyclobutyl, morpholinoethyl, pyrrolidin-1-ylethyl, pyrazol-1-ylethyl or cyclopropylmethyl; wherein A may be optionally substituted on carbon by one or more R⁹;

E is -N(R^a)SO₂-; wherein R^a is hydrogen or methyl;

5 R⁹ is independently selected from hydroxy, methyl, ethenyl, ethynyl, methoxy, ethoxy, propoxy, *N,N*-dimethylamino or methylthio.

In another aspect of the invention, particularly R⁴ is *N*-methylsulphamoyl, *N*-cyclopropylmethylsulphamoyl, *N*-ethylsulphamoyl, *N*-propylsulphamoyl, *N*-allylsulphamoyl, *N*-2-propynylsulphamoyl, *N*-cyclobutylsulphamoyl, *N*-*t*-butylsulphamoyl,
 10 *N*-cyclopropylsulphamoyl, *N*-(2-dimethylaminoethyl)sulphamoyl, *N*-(2-methoxyethyl)sulphamoyl, *N*-(2-pyrazol-1-ylethyl)sulphamoyl, *N*-methyl-*N*-(2-methoxyethyl)sulphamoyl, *N*-(1-cyclopropylethyl)sulphamoyl, *N*-(3-dimethylaminopropyl)sulphamoyl, *N*-(3-methoxypropyl)sulphamoyl, *N*-(3-hydroxy-2-hydroxymethylprop-2-yl)sulphamoyl, *N*-(1,3-dihydroxyprop-2-yl)sulphamoyl,
 15 *N*-(3-morpholino-2-methylprop-2-yl)sulphamoyl, *N*-(1,3-dimethoxyprop-2-yl)sulphamoyl, *N*-(1,3-diethoxyprop-2-yl)sulphamoyl, *N*-(1-methoxyprop-2-yl)sulphamoyl, *N*-(1-ethoxyprop-2-yl)sulphamoyl, *N*-(1-hydroxyprop-2-yl)sulphamoyl, *N*-(3-methylthio-2-methylprop-2-yl)sulphamoyl, *N*-(3-pyrrolidin-1-yl-2-methylprop-2-yl)sulphamoyl,
 20 *N*-(3-methoxy-2-methylprop-2-yl)sulphamoyl, *N*-(2-methoxy-2-methylpropyl)sulphamoyl, *N*-(1-propoxyprop-2-yl)sulphamoyl or *N*-(3-hydroxy-2-methylprop-2-yl)sulphamoyl.

Preferably *q* is 0 or 1.

Preferably *p* + *q* is 1 or 2.

More preferably *p* + *q* is 1.

25 Particularly *p* + *q* is 1 and the R³ or R⁴ group is in the para position of the anilino group.

In another aspect of the invention preferably *p* + *q* is 0, 1 or 2; wherein the values of R³ may be the same or different and the values of R⁴ may be the same or different.

Therefore in a further aspect of the invention, there is provided a compound of formula
 30 (I) (as depicted above) wherein:

Ring A is a group of formula (IA);

one of X¹, X², X³ and X⁴ are nitrogen and the others are CH;

n is 0;

R^2 is hydrogen;

R^3 is sulphonamoyl;

p is 0 or 1;

R^4 is *N*-(2-dimethylaminoethyl)sulphonamoyl, *N*-(2-methoxyethyl)sulphonamoyl, *N*-methylsulphonamoyl or *N*-(3-dimethylaminopropyl)sulphonamoyl.

q is 0 or 1;

$p + q$ is 1 and the R^3 or R^4 group is in the para position of the anilino group;
or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

Therefore in another aspect of the invention, there is provided a compound of formula
10 (I) (as depicted above) wherein:

Ring A is a group of formula (IA) (as depicted above)

wherein one of X^1 , X^2 , X^3 and X^4 is nitrogen and the others are CR^5 ; wherein the values of R^5 may be the same or different;

n is 0;

15 R^2 is selected from hydrogen, C_{1-6} alkyl or *N,N*-(C_{1-6} alkyl)₂amino;

R^5 is selected from hydrogen, C_{1-6} alkyl or C_{1-6} alkoxy; wherein R^5 may be optionally substituted on carbon by one or more R^7 ; wherein R^7 is selected from halo;

R^3 is sulphonamoyl or halo;

p is 0 or 1;

20 R^4 is a group A-E-; wherein

A is selected from C_{1-6} alkyl, C_{3-8} cycloalkyl, (heterocyclic group) C_{1-6} alkyl or C_{3-8} cycloalkyl C_{1-6} alkyl; wherein A may be optionally substituted on carbon by one or more R^9 ;

E is -N(R^a)SO₂-; wherein R^a is hydrogen or C_{1-6} alkyl;

25 R^9 is independently selected from hydroxy, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, *N,N*-(C_{1-6} alkyl)₂amino or C_{1-6} alkylS(O)_a wherein a is 0;

q is 0 or 1;

$p + q$ is 0, 1 or 2; wherein the values of R^3 may be the same or different and the values of R^4 may be the same or different;

30 or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

Therefore in an additional aspect of the invention, there is provided a compound of formula (I) (as depicted above) wherein:

Ring A is a group of formula (IA) (as depicted above)

wherein one of X¹, X², X³ and X⁴ is nitrogen and the others are CR⁵; wherein the values of R⁵ may be the same or different;

n is 0;

5 R² is selected from hydrogen, methyl, ethyl or *N,N*-dimethylamino;

R⁵ is selected from hydrogen, methyl, methoxy or 2,2,2-trifluoroethoxy;

R³ is sulphamoyl or fluoro;

p is 0 or 1;

10 R⁴ is *N*-methylsulphamoyl, *N*-cyclopropylmethylsulphamoyl, *N*-ethylsulphamoyl, *N*-propylsulphamoyl, *N*-allylsulphamoyl, *N*-2-propynylsulphamoyl, *N*-cyclobutylsulphamoyl, *N*-*t*-butylsulphamoyl, *N*-cyclopropylsulphamoyl, *N*-(2-dimethylaminoethyl)sulphamoyl, *N*-(2-methoxyethyl)sulphamoyl, *N*-(2-pyrazol-1-ylethyl)sulphamoyl, *N*-methyl-*N*-(2-methoxyethyl)sulphamoyl, *N*-(1-cyclopropylethyl)sulphamoyl, *N*-(3-dimethylaminopropyl)sulphamoyl, *N*-(3-methoxypropyl)sulphamoyl, *N*-(3-hydroxy-2-hydroxymethylprop-2-yl)sulphamoyl, *N*-(1,3-dihydroxyprop-2-yl)sulphamoyl, *N*-(3-morpholino-2-methylprop-2-yl)sulphamoyl, *N*-(1,3-dimethoxyprop-2-yl)sulphamoyl, *N*-(1,3-diethoxyprop-2-yl)sulphamoyl, *N*-(1-methoxyprop-2-yl)sulphamoyl, *N*-(1-ethoxyprop-2-yl)sulphamoyl, *N*-(1-hydroxyprop-2-yl)sulphamoyl, *N*-(3-methylthio-2-methylprop-2-yl)sulphamoyl, *N*-(3-pyrrolidin-1-yl-2-methylprop-2-yl)sulphamoyl, *N*-(3-methoxy-2-methylprop-2-yl)sulphamoyl, *N*-(2-methoxy-2-methylpropyl)sulphamoyl, *N*-(1-propoxyprop-2-yl)sulphamoyl or *N*-(3-hydroxy-2-methylprop-2-yl)sulphamoyl;

20 q is 0 or 1;

25 p + q is 0, 1 or 2; wherein the values of R³ may be the same or different and the values of R⁴ may be the same or different;

or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.


In another aspect of the invention, preferred compounds of the invention are any one of the Examples or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

30 In another aspect of the invention, preferred compounds of the invention are any one of Examples 7, 27, 29, 39, 41, 42, 46, 63, 64 or 66 or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

Preferred aspects of the invention are those which relate to the compound of formula (I) or a pharmaceutically acceptable salt thereof.


Another aspect of the present invention provides a process for preparing a compound of formula (I) or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof which process (wherein variable groups are, unless otherwise specified, as defined in formula (I)) comprises of:

- 5 a) reaction of a pyrimidine of formula (II):



(II)


wherein L is a displaceable group; with an amine of formula (III):



(III)


10

- b) reacting a pyrimidine of formula (IV):



(IV)

with a compound of the formula (V):




(V)

15

wherein one of M and Q is a displaceable group X and the other is an metallic reagent Y; or


- c) reacting a compounds of formula (VI):

- 13 -



(VI)

with a compound of formula (VII):




(VII)

5


wherein R^5 is C_{1-6} alkyl and R^6 is hydrogen or R^1 ;

d) reacting an amino compound of formula (VIII):



(VIII)

10 with a compound of formula (IX):



wherein E is a displaceable group;

and thereafter if necessary:

i) converting a compound of the formula (I) into another compound of the formula (I);

15 ii) removing any protecting groups;

iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

L is a displaceable group, suitable values for L are for example, a halogeno, methylthio or sulphonyloxy group, for example a chloro, bromo, methanesulphonyloxy or

toluene-4-sulphonyloxy group.

E is a displaceable group, suitable values for E are for example, a halogeno or sulphonyloxy group, for example a chloro, bromo, iodo or trifluoromethanesulphonyloxy group.


- 5 A suitable displaceable group X is, for example, a halogeno or sulphonyl group, for example a bromo, iodo or trifluoromethylsulphonyl group.

A suitable metallic group Y, is, for example, copper, lithium, an organoboron reagent such as $-B(OH)_2$, $-B(OPr^i)_2$ or $-B(Et)_2$, or an organotin compound such as $SnBu_3$, an organosilicon compound such as $Si(Me)F_2$, an organozirconium compound such as $ZrCl_3$, an
 10 organoaluminium compound such as $AlEt_2$, an organomagnesium compound such as $MgBr$, an organozinc compound such as $ZnCl$ or an organomercury compound such as $HgBr$.

Specific reaction conditions for the above reactions are as follows.

- a) Pyrimidines of formula (II) and amines of formula (III) may be reacted together:
 i) in the presence of a suitable solvent for example a ketone such as acetone or an alcohol such
 15 as ethanol or butanol or an aromatic hydrocarbon such as toluene or *N*-methyl pyrrolidine, optionally in the presence of a suitable acid for example an inorganic acid such as hydrochloric acid or sulphuric acid, or an organic acid such as acetic acid or formic acid (or a suitable Lewis acid) and at a temperature in the range of $0^\circ C$ to reflux, preferably reflux; or
 ii) under standard Buchwald conditions (for example see *J. Am. Chem. Soc.*, 118, 7215; *J. Am.*
 20 *Chem. Soc.*, 119, 8451; *J. Org. Chem.*, 62, 1568 and 6066) for example in the presence of palladium acetate, in a suitable solvent for example an aromatic solvent such as toluene, benzene or xylene, with a suitable base for example an inorganic base such as caesium carbonate or an organic base such as potassium-*t*-butoxide, in the presence of a suitable ligand such as 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl and at a temperature in the range of 25 to
 25 $80^\circ C$.

Pyrimidines of the formula (II) may be prepared according to the following scheme:




wherein one of M and Q is a displaceable group X as defined above and the other is an
 metallic reagent Y as defined above.

- 15 -

Cross coupling conditions are well known in the art. Suitable conditions include, for example, those described under b) below.


Where Ring A is a group of formula (IA) and L is methylthio, compounds of the formula (II) may also be prepared according to the following scheme:



5


K is a suitable leaving group (for example C_{1-6} alkanoyloxy), R^5 and R^6 are as defined above, Q is a suitable leaving group (for example C_{1-6} alkoxy).

Where Ring A is a group of formula (IB) and L is methylthio, compounds of the formula (II) may also be prepared according to the following scheme:



wherein R^5 and R^6 are as defined above.

Compounds of formula (IIg) or (III) may be further modified to produce compounds of formula (IIIn):




It will be appreciated by those skilled in the art that compounds of formula (IIIn) may be additionally modified by standard functional group modification reactions known in the art to produce compounds of formula (II) where L is other leaving groups for example chloro, bromo, tosyl and mesyl.

Compounds of formula (IIa), (IIb), (IIc), (II d), (III) and (IIj) are commercially available compounds, or they are known in the literature, or they are prepared by standard processes known in the art.

- b) Compounds of formula (IV) and compounds of formula (V) may be reacted together under standard cross coupling conditions. Examples of these are in the presence of a catalyst, for example, a metallic catalyst such as tetrakis(triphenylphosphine)palladium(0), palladium(II) chloride, palladium(II) bromide, nickel(II) chloride, nickel(II) bromide or bis(triphenylphosphine)nickel(II) chloride, in the presence of a suitable inert solvent or diluent, for example tetrahydrofuran, 1,4-dioxan, 1,2-dimethoxyethane, benzene, toluene, xylene, methanol or ethanol. The reaction is preferably conducted in the presence of a suitable base such as, for example, sodium carbonate or potassium carbonate, pyridine, 4-dimethylaminopyridine, triethylamine or morpholine, and conveniently at a temperature in the range, for example 10 to 250°C, preferably in the range 60 to 120°C.

Compounds of formula (IV) may be prepared according to the following scheme:



Compounds of formula (V) are commercially available compounds, or they are known in the literature, or they are prepared by standard processes known in the art.

- c) compounds of formula (VI) and compounds of formula (VII) are reacted together in a suitable solvent such as *N*-methylpyrrolidinone or butanol at a temperature in the range of 100-200°C, preferably in the range of 150-170°C. The reaction is preferably conducted in the presence of a suitable base such as, for example, sodium methoxide or potassium carbonate.

Compounds of formula (VI) and (VII) are commercially available compounds, or they are known in the literature, or they are prepared by standard processes known in the art, or compounds of formula (VII) may be prepared by a process similar to that described for (III f) and (III k) hereinabove.

- d) Amino compounds of formula (VIII) and compounds of formula (IX) may be reacted together under standard Buchwald conditions such as those described under process a(ii) above.

Compounds of formula (VIII) are prepared as described above.

- Compounds of formula (IX) are commercially available compounds, or they are known in the literature, or they are prepared by standard processes known in the art.

It will be appreciated that certain of the various ring substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halogeno group. Particular examples of modifications include the reduction of a nitro group to an amino group by for example, catalytic hydrogenation with a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating; oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in the compounds. The instances where protection is necessary or desirable and suitable methods for protection are known to those skilled in the art. Conventional protecting groups may be used in accordance with standard practice (for illustration see T.W. Green, Protective Groups in Organic Synthesis, John Wiley and Sons, 1991). Thus, if reactants include groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or *t*-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an

arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art.

As stated hereinbefore the compounds defined in the present invention possesses anti-cell-proliferation activity such as anti-cancer activity which is believed to arise from the CDK inhibitory activity of the compound. These properties may be assessed, for example, using the procedure set out below:-

Assay

The following abbreviations have been used :-

HEPES is *N*-[2-Hydroxyethyl]piperazine-*N'*-[2-ethanesulfonic acid]

DTT is Dithiothreitol

PMSF is Phenylmethylsulfonyl fluoride

The compounds were tested in an *in vitro* kinase assay in 96 well format using Scintillation Proximity Assay (SPA - obtained from Amersham) for measuring incorporation

of [γ -33-P]-Adenosine Triphosphate into a test substrate (GST-Retinoblastoma protein; GST-Rb). In each well was placed the compound to be tested (diluted in DMSO and water to correct concentrations) and in control wells either roscovitine as an inhibitor control or DMSO as a positive control.

5 Approximately 0.2 μ l of CDK2/Cyclin E partially-purified enzyme (amount dependent on enzyme activity) diluted in 25 μ l incubation buffer was added to each well then 20 μ l of GST-Rb/ATP/ATP33 mixture (containing 0.5 μ g GST-Rb and 0.2 μ M ATP and 0.14 μ Ci [γ -33-P]-Adenosine Triphosphate in incubation buffer), and the resulting mixture shaken gently, then incubated at room temperature for 60 minutes.

10 To each well was then added 150 μ L stop solution containing (0.8mg/well of Protein A-PVT SPA bead (Amersham)), 20pM/well of Anti-Glutathione Transferase, Rabbit IgG (obtained from Molecular Probes), 61mM EDTA and 50mM HEPES pH 7.5 containing 0.05% sodium azide.

 The plates were sealed with Topseal-S plate sealers, left for two hours then spun at
15 2500rpm, 1124xg., for 5 minutes. The plates were read on a Topcount for 30 seconds per well.

 The incubation buffer used to dilute the enzyme and substrate mixes contained 50mM HEPES pH7.5, 10mM MnCl₂, 1mM DTT, 100 μ M Sodium vanadate, 100 μ M NaF, 10mM Sodium Glycerophosphate, BSA (1mg/ml final).

Test substrate

20 In this assay only part of the retinoblastoma protein (Science 1987 Mar13;235(4794):1394-1399; Lee W.H., Bookstein R., Hong F., Young L.J., Shew J.Y., Lee E.Y.) was used, fused to a GST tag. PCR of retinoblastoma gene encoding amino acids 379-928 (obtained from retinoblastoma plasmid ATCC pLRbRNL) was performed, and the sequence cloned into pGEX 2T fusion vector (Smith D.B. and Johnson, K.S. Gene 67, 31
25 (1988); which contained a tac promoter for inducible expression, internal lac I^q gene for use in any E.Coli host, and a coding region for thrombin cleavage - obtained from Pharmacia Biotech) which was used to amplify amino acids 792-928. This sequence was again cloned into pGEX 2T.

 The retinoblastoma 792-928 sequence so obtained was expressed in E.Coli (BL21
30 (DE3) pLysS cells) using standard inducible expression techniques, and purified as follows.

 E.coli paste was resuspended in 10ml/g of NETN buffer (50mM Tris pH 7.5, 120mM NaCl, 1mM EDTA, 0.5%v/v NP-40, 1mM PMSF, 1 μ g/ml leupeptin, 1 μ g/ml aprotinin and

1ug/ml pepstatin) and sonicated for 2 x 45 seconds per 100ml homogenate. After centrifugation, the supernatant was loaded onto a 10ml glutathione Sepharose column (Pharmacia Biotech, Herts, UK), and washed with NETN buffer. After washing with kinase buffer (50mM HEPES pH 7.5, 10mM MgCl₂, 1mM DTT, 1mM PMSF, 1ug/ml leupeptin, 1ug/ml aprotinin and 1ug/ml pepstatin) the protein was eluted with 50mM reduced glutathione in kinase buffer. Fractions containing GST-Rb(792-927) were pooled and dialysed overnight against kinase buffer. The final product was analysed by Sodium Dodeca Sulfate (SDS) PAGE (Polyacrylamide gel) using 8-16% Tris-Glycine gels (Novex, San Diego, USA).

CDK2 and Cyclin E

- 10 The open reading frames of CDK2 and Cyclin E were isolated by reverse transcriptase-PCR using HeLa cell and activated T cell mRNA as a template and cloned into the insect expression vector pVL1393 (obtained from Invitrogen 1995 catalogue number: V1392-20). CDK2 and cyclin E were then dually expressed [using a standard virus Baculogold co-infection technique] in the insect SF21 cell system (Spodoptera Frugiperda cells derived from ovarian tissue of the Fall Army Worm - commercially available).

Example production of Cyclin E/CDK2

The following Example provides details of the production of Cyclin E/CDK2 in SF21 cells (in TC100 + 10% FBS(TCS) + 0.2% Pluronic) having dual infection MOI 3 for each virus of Cyclin E & CDK2.

- 20 SF21 cells grown in a roller bottle culture to 2.33×10^6 cells/ml were used to inoculate 10 x 500 ml roller bottles at 0.2×10^6 cells/ml. The roller bottles were incubated on a roller rig at 28°C.

- After 3 days (72 hrs.) the cells were counted, and the average from 2 bottles found to be 1.86×10^6 cells/ml. (99% viable). The cultures were then infected with the dual viruses at an MOI 3 for each virus.

The viruses were mixed together before addition to the cultures, and the cultures returned to the roller rig 28°C.

- After 2 days (48 hrs.) post infection the 5 Litres of culture was harvested. The total cell count at harvest was 1.58×10^6 cells/ml.(99% viable). The cells were spun out at 2500rpm, 30 mins., 4°C in Heraeus Omnifuge 2.0 RS in 250 ml. lots. The supernatant was discarded.

Partial co-purification of Cdk2 and Cyclin E

Sf21 cells were resuspended in lysis buffer (50mM Tris pH 8.2, 10mM MgCl₂, 1mM DTT, 10mM glycerophosphate, 0.1mM sodium orthovanadate, 0.1mM NaF, 1mM PMSF, 1µg/ml leupeptin and 1µg/ml aprotinin) and homogenised for 2 minutes in a 10ml Dounce homogeniser. After centrifugation, the supernatant was loaded onto a Poros HQ/M 1.4/100 anion exchange column (PE Biosystems, Hertford, UK). Cdk2 and Cyclin E were coeluted at the beginning of a 0-1M NaCl gradient (run in lysis buffer minus protease inhibitors) over 20 column volumes. Co-elution was checked by western blot using both anti-Cdk2 and anti-Cyclin E antibodies (Santa Cruz Biotechnology, California, US).

By analogy, assays designed to assess inhibition of CDK4 and CDK6 may be constructed. CDK2 (EMBL Accession No. X62071) may be used together with Cyclin A or Cyclin E (see EMBL Accession No. M73812), and further details for such assays are contained in PCT International Publication No. WO99/21845, the relevant Biochemical & Biological Evaluation sections of which are hereby incorporated by reference.

Although the pharmacological properties of the compounds of the formula (I) vary with structural change, in general activity possessed by compounds of the formula (I) may be demonstrated at IC₅₀ concentrations or doses in the range 250µM to 1nM.

The *in vivo* activity of the compounds of the present invention may be assessed by standard techniques, for example by measuring inhibition of cell growth and assessing cytotoxicity.

Inhibition of cell growth may be measured by staining cells with Sulforhodamine B (SRB), a fluorescent dye that stains proteins and therefore gives an estimation of amount of protein (i.e. cells) in a well (see Boyd, M.R.(1989) Status of the NCI preclinical antitumour drug discovery screen. *Prin. Prac Oncol* 10:1-12). Thus, the following details are provided of measuring inhibition of cell growth :-

Cells were plated in appropriate medium in a volume of 100µl in 96 well plates; media was Dulbecco's Modified Eagle media for MCF-7, SK-UT-1B and SK-UT-1. The cells were allowed to attach overnight, then inhibitor compounds were added at various concentrations in a maximum concentration of 1% DMSO (v/v). A control plate was assayed to give a value for cells before dosing. Cells were incubated at 37°C, (5% CO₂) for three days.

At the end of three days TCA was added to the plates to a final concentration of 16% (v/v). Plates were then incubated at 4°C for 1 hour, the supernatant removed and the plates washed in tap water. After drying, 100µl SRB dye (0.4% SRB in 1% acetic acid) was added

for 30 minutes at 37°C. Excess SRB was removed and the plates washed in 1% acetic acid. The SRB bound to protein was solubilised in 10mM Tris pH7.5 and shaken for 30 minutes at room temperature. The ODs were read at 540nm, and the concentration of inhibitor causing 50% inhibition of growth was determined from a semi-log plot of inhibitor concentration versus absorbance. The concentration of compound that reduced the optical density to below that obtained when the cells were plated at the start of the experiment gave the value for toxicity.

Typical IC₅₀ values for compounds of the invention when tested in the SRB assay are in the range 1mM to 1nM.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore in association with a pharmaceutically-acceptable diluent or carrier.

The composition may be in a form suitable for oral administration, for example as a tablet or capsule, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) as a sterile solution, suspension or emulsion, for topical administration as an ointment or cream or for rectal administration as a suppository.

In general the above compositions may be prepared in a conventional manner using conventional excipients.

The compound of formula (I) will normally be administered to a warm-blooded animal at a unit dose within the range 5-5000 mg per square meter body area of the animal, i.e. approximately 0.1-100 mg/kg, and this normally provides a therapeutically-effective dose. A unit dose form such as a tablet or capsule will usually contain, for example 1-250 mg of active ingredient. Preferably a daily dose in the range of 1-50 mg/kg is employed. However the daily dose will necessarily be varied depending upon the host treated, the particular route of administration, and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any particular patient.

According to a further aspect of the present invention there is provided a compound of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore for use in a method of treatment of the human or animal body by therapy.

We have found that the compounds defined in the present invention, or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, are effective cell cycle

inhibitors (anti-cell proliferation agents), which property is believed to arise from their CDK inhibitory properties. Accordingly the compounds of the present invention are expected to be useful in the treatment of diseases or medical conditions mediated alone or in part by CDK enzymes, i.e. the compounds may be used to produce a CDK inhibitory effect in a

5 warm-blooded animal in need of such treatment. Thus the compounds of the present invention provide a method for treating the proliferation of malignant cells characterised by inhibition of CDK enzymes, i.e. the compounds may be used to produce an anti-proliferative effect mediated alone or in part by the inhibition of CDKs. Such a compound of the invention is expected to possess a wide range of anti-cancer properties as CDKs have been implicated in

10 many common human cancers such as leukaemia and breast, lung, colon, rectal, stomach, prostate, bladder, pancreas and ovarian cancer. Thus it is expected that a compound of the invention will possess anti-cancer activity against these cancers. It is in addition expected that a compound of the present invention will possess activity against a range of leukaemias, lymphoid malignancies and solid tumours such as carcinomas and sarcomas in tissues such as

15 the liver, kidney, prostate and pancreas. In particular such compounds of the invention are expected to slow advantageously the growth of primary and recurrent solid tumours of, for example, the colon, breast, prostate, lungs and skin. More particularly such compounds of the invention, or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, are expected to inhibit the growth of those primary and recurrent solid tumours which are

20 associated with CDKs, especially those tumours which are significantly dependent on CDKs for their growth and spread, including for example, certain tumours of the colon, breast, prostate, lung, vulva and skin.

It is further expected that a compound of the present invention will possess activity against other cell-proliferation diseases in a wide range of other disease states including

25 leukaemias, fibroproliferative and differentiative disorders, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, atherosclerosis, arterial restenosis, autoimmune diseases, acute and chronic inflammation, bone diseases and ocular diseases with retinal vessel proliferation.

Thus according to this aspect of the invention there is provided a compound of the

30 formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore for use as a medicament; and the use of a compound of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore in the manufacture of a medicament for use in the production of a cell cycle

inhibitory (anti-cell-proliferation) effect in a warm-blooded animal such as man. Particularly, an inhibitory effect is produced by preventing entry into or progression through the S phase by inhibition of CDK2, CDK4 and/or CDK6, especially CDK2.

According to a further feature of the invention there is provided the use of a compound
5 of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore in the manufacture of a medicament for use in the treatment of cancers (solid tumours and leukemias), fibroproliferative and differentiative disorders, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, atherosclerosis, arterial restenosis, autoimmune diseases, acute and chronic
10 inflammation, bone diseases and ocular diseases with retinal vessel proliferation, particularly in the treatment of cancers.

According to a further feature of this aspect of the invention there is provided a method for producing a cell cycle inhibitory (anti-cell-proliferation) effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal
15 an effective amount of a compound as defined immediately above. Particularly, an inhibitory effect is produced by preventing entry into or progression through the S phase by inhibition of CDK2, CDK4 and/or CDK6, especially CDK2.

According to an additional feature of this aspect of the invention there is provided a method of treating cancers (solid tumours and leukaemias), fibroproliferative and
20 differentiative disorders, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, atherosclerosis, arterial restenosis, autoimmune diseases, acute and chronic inflammation, bone diseases and ocular diseases with retinal vessel proliferation, in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I) or a
25 pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof as defined hereinbefore.

Particularly there is provided a method of treating cancer in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof as defined hereinbefore.

30 In a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore in association with a

pharmaceutically-acceptable diluent or carrier for use in the production of a cell cycle inhibitory (anti-cell-proliferation) effect in a warm-blooded animal such as man.

In a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore in association with a pharmaceutically-acceptable diluent or carrier for use in the treatment of cancers (solid tumours and leukaemias), fibroproliferative and differentiative disorders, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, atherosclerosis, arterial restenosis, autoimmune diseases, acute and chronic inflammation, bone diseases and ocular diseases with retinal vessel proliferation, in a warm-blooded animal such as man.

In a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore in association with a pharmaceutically-acceptable diluent or carrier for use in the treatment of cancer in a warm-blooded animal such as man.

In a further aspect of the invention there is provided a compound of formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore, for use in the production of a cell cycle inhibitory (anti-cell-proliferation) effect in a warm-blooded animal such as man.

In a further aspect of the invention there is provided a compound of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore for use in the treatment of cancers (solid tumours and leukaemias), fibroproliferative and differentiative disorders, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, atherosclerosis, arterial restenosis, autoimmune diseases, acute and chronic inflammation, bone diseases and ocular diseases with retinal vessel proliferation, in a warm-blooded animal such as man.

In a further aspect of the invention there is provided a compound of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore for use in the treatment of cancer in a warm-blooded animal such as man.

Preventing cells from entering DNA synthesis by inhibition of essential S-phase initiating activities such as CDK2 initiation may also be useful in protecting normal cells of

the body from toxicity of cycle-specific pharmaceutical agents. Inhibition of CDK2 or 4 will prevent progression into the cell cycle in normal cells which could limit the toxicity of cycle-specific pharmaceutical agents which act in S-phase, G2 or mitosis. Such protection may result in the prevention of hair loss normally associated with these agents.

5 Therefore in a further aspect of the invention there is provided a compound of formula (I) as defined above or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof for use as a cell protective agent.

Therefore in a further aspect of the invention there is provided a compound of formula (I) as defined above or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof
10 for use in preventing hair loss arising from the treatment of malignant conditions with pharmaceutical agents.

Examples of pharmaceutical agents for treating malignant conditions that are known to cause hair loss include alkylating agents such as ifosfamide and cyclophosphamide; antimetabolites such as methotrexate, 5-fluorouracil, gemcitabine and cytarabine; vinca
15 alkaloids and analogues such as vincristine, vinblastine, vindesine, vinorelbine; taxanes such as paclitaxel and docetaxel; topoisomerase I inhibitors such as irinotecan and topotecan; cytotoxic antibiotics such as doxorubicin, daunorubicin, mitoxantrone, actinomycin-D and mitomycin; and others such as etoposide and tretinoin.

In another aspect of the invention, the compound of formula (I), or a pharmaceutically
20 acceptable salt or *in vivo* hydrolysable ester thereof, may be administered in association with a one or more of the above pharmaceutical agents. In this instance the compound of formula (I) may be administered by systemic or non systemic means. Particularly the compound of formula (I) may be administered by non-systemic means, for example topical administration.

Therefore in an additional feature of the invention, there is provided a method of
25 preventing hair loss during treatment for one or more malignant conditions with pharmaceutical agents, in a warm-blooded animal, such as man, which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

In an additional feature of the invention, there is provided a method of preventing hair
30 loss during treatment for one or more malignant conditions with pharmaceutical agents, in a warm-blooded animal, such as man, which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt or *in vivo*

vivo hydrolysable ester thereof in simultaneous, sequential or separate administration with an effective amount of said pharmaceutical agent.

According to a further aspect of the invention there is provided a pharmaceutical composition for use in preventing hair loss arising from the treatment of malignant conditions with pharmaceutical agents which comprises a compound of formula (I), or a
5 pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, and said pharmaceutical agent, in association with a pharmaceutically acceptable diluent or carrier.

According to a further aspect of the present invention there is provided a kit comprising a compound of formula (I), or a pharmaceutically acceptable salt or *in vivo*
10 hydrolysable ester thereof, and a pharmaceutical agent for treating malignant conditions that is known to cause hair loss.

According to a further aspect of the present invention there is provided a kit comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable
15 ester thereof, in a first unit dosage form;
- b) a pharmaceutical agent for treating malignant conditions that is known to cause hair loss; in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

According to another feature of the invention there is provided the use of a compound
20 of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, in the manufacture of a medicament for the prevention of hair loss during treatment of malignant conditions with pharmaceutical agents.

According to a further aspect of the present invention there is provided a combination treatment for the prevention of hair loss comprising the administration of an effective amount
25 of a compound of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the simultaneous, sequential or separate administration of an effective amount of a pharmaceutical agent for treatment of malignant conditions to a warm-blooded animal, such as man.

30 As stated above the size of the dose required for the therapeutic or prophylactic treatment of a particular cell-proliferation disease will necessarily be varied depending on the host treated, the route of administration and the severity of the illness being treated. A unit

dose in the range, for example, 1-100 mg/kg, preferably 1-50 mg/kg is envisaged.

The CDK inhibitory activity defined hereinbefore may be applied as a sole therapy or may involve, in addition to a compound of the invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous, sequential
5 or separate administration of the individual components of the treatment. In the field of medical oncology it is normal practice to use a combination of different forms of treatment to treat each patient with cancer. In medical oncology the other component(s) of such conjoint treatment in addition to the cell cycle inhibitory treatment defined hereinbefore may be: surgery, radiotherapy or chemotherapy. Such chemotherapy may cover three main categories
10 of therapeutic agent:

(i) other cell cycle inhibitory agents that work by the same or different mechanisms from those defined hereinbefore;

(ii) cytostatic agents such as antioestrogens (for example tamoxifen, toremifene, raloxifene, droloxifene, idoxifene), progestogens (for example megestrol acetate), aromatase
15 inhibitors (for example anastrozole, letrozole, vorazole, exemestane), antiprogestogens, antiandrogens (for example flutamide, nilutamide, bicalutamide, cyproterone acetate), LHRH agonists and antagonists (for example goserelin acetate, luproline), inhibitors of testosterone 5 α -dihydroreductase (for example finasteride), anti-invasion agents (for example metalloproteinase inhibitors like marimastat and inhibitors of urokinase plasminogen activator
20 receptor function) and inhibitors of growth factor function, (such growth factors include for example platelet derived growth factor and hepatocyte growth factor such inhibitors include growth factor antibodies, growth factor receptor antibodies, tyrosine kinase inhibitors and serine/threonine kinase inhibitors); and

(iii) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical
25 oncology, such as antimetabolites (for example antifolates like methotrexate, fluoropyrimidines like 5-fluorouracil, purine and adenosine analogues, cytosine arabinoside); antitumour antibiotics (for example anthracyclines like doxorubicin, daunomycin, epirubicin and idarubicin, mitomycin-C, dactinomycin, mithramycin); platinum derivatives (for example cisplatin, carboplatin); alkylating agents (for example nitrogen mustard, melphalan,
30 chlorambucil, busulphan, cyclophosphamide, ifosfamide, nitrosoureas, thiotepa); antimetotic agents (for example vinca alkaloids like vincristine and taxoids like taxol, taxotere); topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide,

amsacrine, topotecan). According to this aspect of the invention there is provided a pharmaceutical product comprising a compound of the formula (I) as defined hereinbefore and an additional anti-tumour substance as defined hereinbefore for the conjoint treatment of cancer.

5 In addition to their use in therapeutic medicine, the compounds of formula (I) and their pharmaceutically acceptable salts are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effects of inhibitors of cell cycle activity in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

10 In the above other pharmaceutical composition, process, method, use and medicament manufacture features, the alternative and preferred embodiments of the compounds of the invention described herein also apply.

Examples

The invention will now be illustrated by the following non-limiting examples in which, unless stated otherwise:

- (i) temperatures are given in degrees Celsius (°C); operations were carried out at room or ambient temperature, that is, at a temperature in the range of 18-25°C;
- (ii) organic solutions were dried over anhydrous magnesium sulphate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 Pascals;
- 20 4.5-30mmHg) with a bath temperature of up to 60°C;
- (iii) chromatography means flash chromatography on silica gel; thin layer chromatography (TLC) was carried out on silica gel plates; where an amino isolate column is referred to, this means a column containing 10g of NH₂ SPE sorbent of 40 micron particle size, the sorbent being contained in a 60ml disposable syringe and supported by a porous disc, obtained from
- 25 IST, Mid Glamorgan, UK under the name "Isolute NH₂", "Isolute" is a trademark;
- (iv) in general, the course of reactions was followed by TLC and reaction times are given for illustration only;
- (v) final products had satisfactory proton nuclear magnetic resonance (NMR) spectra and/or mass spectral data;
- 30 (vi) yields are given for illustration only and are not necessarily those which can be obtained by diligent process development; preparations were repeated if more material was required;
- (vii) when given, NMR data is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal standard,

determined at 300 MHz using perdeuterio dimethyl sulphoxide (DMSO-d₆) as solvent unless otherwise indicated;

(viii) chemical symbols have their usual meanings; SI units and symbols are used;

(ix) solvent ratios are given in volume:volume (v/v) terms; and

5 (x) mass spectra were run with an electron energy of 70 electron volts in the chemical ionization (CI) mode using a direct exposure probe; where indicated ionization was effected by electron impact (EI), fast atom bombardment (FAB) or electrospray (ESP); values for m/z are given; generally, only ions which indicate the parent mass are reported;

(xi) unless stated otherwise compounds containing an asymmetrically substituted carbon
10 and/or sulphur atom have not been resolved;

(xii) where a synthesis is described as being analogous to that described in a previous example the amounts used are the millimolar ratio equivalents to those used in the previous example;

(xvi) the following abbreviations have been used:

	DMF	<i>N,N</i> -dimethylformamide;
15	DMFDMA	<i>N,N</i> -dimethylformamidedimethylacetyl;
	DMSO	dimethylsulphoxide; and
	THF	tetrahydrofuran.

Example 1

20 2-(4-Sulphamoylanilino)-4-(imidazo[1,2a]pyrazin-3-yl)pyrimidine

Thionyl chloride (4ml) and then DMF (5μl) were added to 2-(4-sulphonoylanilino)-4-(imidazo[1,2a]pyrazin-3-yl)pyrimidine (Method 4; 75mg, 0.204mmol) under nitrogen and the mixture was stirred at ambient temperature for 48 hours. The volatiles were removed by evaporation and methanolic ammonia (6ml of a 7M solution) was added directly to the residue
25 under nitrogen and the mixture stirred for 1 hour at ambient temperature. The volatiles were removed by evaporation, and the residue purified by preparative HPLC (on a Gilson, 1" C18 column with gradient elution of 0.01% TFA buffered acetonitrile / water 5% to 50% over 10 minutes and 50% to 95% over 3.5 minutes). Fractions containing product were combined and lyophilised to give the title compound (17 mg, 23%) as a white solid. NMR: 7.20 (s, 2H), 7.60
30 (d, 1H), 7.72 (d, 2H), 7.95 (d, 2H), 8.18 (d, 1H), 8.62 (d, 1H), 8.85 (s, 1H), 9.28 (s, 1H), 10.01(s, 1H), 10.20 (s, 1H); m/z: 368 [MH]⁺.

Example 22-(4-{N-[2-(Dimethylamino)ethyl]sulphamoyl}anilino)-4-(imidazo[1,2a]pyrazin-3-yl)pyrimidine

2-(4-Sulphonoylanilino)-4-(imidazo[1,2a]pyrazin-3-yl)pyrimidine (Method 4; 67mg, 0.181mmol) was dissolved in acetonitrile (1ml) and sulfolane (1ml) and stirred under nitrogen. Phosphoryl chloride (67µl, 0.724mmol) and *N,N*-dimethylacetamide (10µl) were added. The reaction mixture was heated under reflux for 30 minutes, allowed to cool and the volatiles removed by evaporation. The resulting solution was placed under nitrogen and an anhydrous solution of triethylamine (252µl, 1.81mmol) and 1,1-dimethylethyldiamine (40µl, 0.362mmol) in methanol (2ml) was added slowly. The mixture was left at ambient temperature for 30 minutes, then dichloromethane (15ml) and water (15ml) were added. The aqueous layer was separated and adjusted to pH 9 by careful addition of 2M aqueous sodium hydroxide solution. The resulting precipitate was collected by filtration, washed with water (2 x 15ml) and ether (3 x 15ml) and dried over phosphorus pentoxide for 16 hours to give the title compound (16mg, 20%). NMR: 2.1 (s, 6H), 2.3 (t, 2H), 2.8 (t, 2H), 7.3 (br s, 1H), 7.5 (d, 1H), 7.75 (d, 2H), 7.95 (d, 2H), 8.2 (d, 1H), 8.6 (d, 1H), 8.8 (s, 1H), 9.3 (s, 1H), 9.95 (d, 1H), 10.2 (s, 1H); m/z: 440 [MH]⁺.

Example 32-{4-[N-(2-Methoxyethyl)sulphamoyl]anilino}-4-(imidazo[1,2a]pyrazin-3-yl)pyrimidine

Thionyl chloride (6ml) and DMF (10µl) were added to 2-(4-sulphonoylanilino)-4-(imidazo[1,2a]pyrazin-3-yl)pyrimidine (Method 4; 110mg, 0.299mmol) under and the mixture was stirred at ambient temperature for 12 hours. The volatiles were removed by evaporation and the residue placed under nitrogen. Anhydrous pyridine (7ml) followed by 2-methoxyethylamine (26µl, 0.299mmol) was added directly and the mixture stirred at ambient temperature for 3 hours. The volatiles were removed by evaporation, and the residue purified by column chromatography eluting with dichloromethane/methanol (100:0 increasing in polarity to 95:5) to give the title compound (6mg, 5%) as a white solid. NMR: 2.9 (t, 2H), 3.2 (s, 3H), 3.4 (t, 2H), 7.5 (t, 1H), 7.7 (d, 1H), 7.8 (d, 2H), 7.95 (d, 2H), 8.2 (d, 1H), 8.6 (d, 1H), 8.9 (s, 1H), 9.3 (s, 1H), 10.0 (d, 1H), 10.2 (s, 1H); m/z: 426 [MH]⁺.

Example 42-[4-[N-(2-Methoxyethyl)sulphamoyl]anilino]-4-(imidazo[1,2a]pyrimidin-3-yl)pyrimidine

2-[4-Sulphonoylanilino]-4-(imidazo[1,2a]pyrimidin-3-yl)pyrimidine (Method 8; 180mg, 0.49mmol) was dissolved in acetonitrile (1ml) and sulfolane (1ml) and stirred under nitrogen. Phosphoryl chloride (136µl, 1.46mmol) and *N,N*-dimethylacetamide (1.8µl) were added and the reaction mixture was stirred at ambient temperature for 4 hours then heated under reflux for 30 minutes, allowed to cool and the volatiles removed by evaporation. The resulting solution was placed under nitrogen and a solution of 2-methoxyethylamine (85µl, 0.978mmol), triethylamine (204ml, 1.47mmol) in methanol (5ml) was added and the mixture stirred at ambient temperature for 1 hour. The resulting solid was collected by filtration, washed with ether (3 x 20ml), and dried to give the title compound (89mg, 43%). NMR: 2.9 (dd, 1H), 3.08 (m, 2H), 3.18 (s, 3H), 3.3 (t, 2H), 7.56 (d, 1H) 7.77 (d, 2H) 7.98 (d, 2H), 8.6 (d, 1H), 8.86 (dd, 1H), 8.98 (s, 1H), 10.3 (s, 1H) 10.54 (s, 1H); m/z: 426 [MH]⁺.

Example 52-(4-sulphamoylanilino)-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine

2-Anilino-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine (Example 10; 100mg, 0.347mmol) was dissolved in thionyl chloride (2.0ml) and the solution stirred and cooled to 0°C. Chlorosulphonic acid (92µl, 1.39mmol) was added and the reaction was stirred at 0°C for 30 minutes and then at ambient temperature for 1 hour and finally heated at 80°C for 90 minutes. The mixture was allowed to cool and the volatiles were removed by evaporation. 2M methanolic ammonia (6.0ml) was added to the residue in the flask and the mixture stirred at ambient temperature for 1 hour. The volatiles were removed by evaporation and distilled water added to the residue and the mixture stirred for 30 minutes. The precipitate was collected by filtration, washed with water and dried. The crude product was triturated with dichloromethane containing a trace of methanol, collected by filtration washed with dichloromethane / methanol and dried to give the title compound (81mg, 64%) as a white solid. NMR: 7.1 (s, 2H), 7.45 (d, 1H), 7.8 (d, 2H), 8.03 (m, 3H), 8.32 (d, 1H), 8.64 (m, 2H), 8.8 (d, 1H); 10.05 (s, 1H); m/z: 368 [MH]⁺.

Example 62-[4-(N-Methylsulphamoyl)anilino]-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine

2-Anilino-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine (Example 10; 100mg, 0.347mmol) was treated by the method described in Example 5 to give the crude sulphonyl chloride which was treated with 2.0M solution of methylamine in methanol (6.0ml) and the reaction worked-up as described in Example 5 to give the title compound (78mg, 59%) as a white solid. NMR: 2.4 (d, 3H), 7.16 (q, 1H), 7.45 (dd, 1H), 7.75 (d, 2H), 8.07 (m, 3H), 8.33 (d, 1H), 8.67 (m, 2H), 8.8 (d, 1H); 10.11 (s, 1H); m/z: 380 [M-H].

Example 72-{4-[N-(2-Methoxyethyl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine

2-Anilino-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine (Example 10; 100mg, 0.347mmol) was treated by the method described in Example 5 to give the crude sulphonyl chloride which was treated with a solution of 2-methoxyethylamine (0.452ml) in methanol (1.0ml) and the reaction worked-up as described in Example 5 to give the title compound (39mg, 26%) as a white solid. NMR (DMSO d₆ + d₄ Acetic acid): 2.87 (t, 2H), 3.15 (s, 3H), 3.27 (t, 2H), 7.4 (dd, 1H), 7.74 (d, 2H), 8.05 (m, 3H), 8.3 (d, 1H), 8.64 (m, 2H), 8.76 (d, 1H). m/z: 424 [M-H].

Example 82-[4-(N-(3-Dimethylaminopropyl)sulphamoyl)anilino]-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine

2-Anilino-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine (Example 10; 100mg, 0.347mmol) was treated by the method described in Example 5 to give the crude sulphonyl chloride which was treated with a solution of 3-dimethylaminopropylamine (87µl), dimethylethylamine (2.16ml) in methanol (4.0ml) and the resulting suspension stirred at ambient temperature for 90 minutes. Volatiles were removed by evaporation, distilled water was added to the residue and the mixture stirred for 30 minutes. The resulting precipitate was collected by filtration, washed with distilled water and dried. The crude product dissolved in DMF was purified on a 10g amino isolate column pre-equilibrated with methanol. The product was eluted with methanol, the purified product was triturated with ether / ethyl acetate, collected by filtration, washed with ether and dried to the title compound (53mg, 26%) as a

white solid. NMR: 1.48 (m, 2H), 2.03 (s, 6H), 2.13 (t, 2H), 2.75 (t, 2H), 7.33 (s, 1H), 7.45 (dd, 1H), 7.73 (d, 2H), 8.07 (m, 3H), 8.33 (d, 1H), 8.65 (s, 1H), 8.68 (d, 1H), 8.8 (d, 1H), 10.11 (s, 1H); m/z: 451 [M-H]⁻.

5. Example 9

2-Anilino-4-(imidazo[1,2a]pyrazin-3-yl)pyrimidine

3-(3-Dimethylaminoprop-2-en-1-yl)imidazo[1,2a]pyrazine (Method 3; 400mg, 1.85mmol), phenyl guanidine hydrogencarbonate (365mg, 1.85mmol) and sodium methoxide (199mg, 3.7mmol) were dissolved in anhydrous dimethylacetamide (15ml) and the mixture heated under reflux at 160°C under nitrogen for 2 hours. The reaction mixture was allowed to cool, acetic acid (212μl, 3.7mmol) was added and the mixture poured carefully on to ice-water (40ml). The ice was allowed to melt and the aqueous solution extracted with dichloromethane (2 x 20ml). The extracts were combined, dried and the solvent removed by evaporation to give title compound (392mg, 72%) as a brown solid. NMR: 7.0 (t, 1H), 7.35 (t, 2H), 7.48 (d, 1H), 7.75 (d, 2H), 8.1 (d, 1H), 8.52 (d, 1H), 8.88 (s, 1H), 9.21 (s, 1H), 9.86 (s, 1H), 9.98 (d, 1H); m/z: 299 [MH]⁺.

Example 10

2-Anilino-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine

3-(3-Dimethylaminoprop-2-en-1-yl)imidazo[1,2b]pyridazine (Method 17; 600mg, 2.78mmol), phenyl guanidine hydrogen carbonate (603mg, 3.06mmol) and sodium methoxide (300mg, 5.56mmol) were suspended in dry dimethyl acetamide (25ml) under nitrogen and the mixture was stirred and heated at 160°C for 2 hours. The reaction was then stirred at room temperature for 20 hours and then further phenyl guanidine hydrogen carbonate (273mg) and sodium methoxide (150mg) were added and the reaction heated at 160°C for 2 hours. The reaction was allowed to cool to ambient temperature and acetic acid (0.5ml) added and the volatiles removed by evaporation. Water was added to the residue and the resulting suspension stirred for 30 minutes. The product was collected by filtration, washed with water and dried to give the title compound (705mg, 88%) as a brown solid. NMR: 6.96 (t, 1H), 7.33 (m, 2H), 7.42 (dd, 1H), 7.85 (d, 2H), 7.97 (d, 1H), 8.30 (d, 1H), 8.55 (s, 1H), 8.61 (d, 1H), 8.77 (d, 1H), 9.63 (s, 1H). m/z: 289 [MH]⁺.

Example 11**2-Anilino-4-(imidazo[1,2a]pyrimidin-3-yl)pyrimidine**

3-(3-Dimethylaminoprop-2-en-1-yl)imidazo[1,2a]pyrimidine (Method 7; 450mg, 2.08mmol) was treated as described in Example 9 to give crude product which was triturated with ether to give the title compound as the major component (70%) in an isomeric mixture with 2-anilino-4-(imidazo[1,2a]pyrimidin-2-yl)pyrimidine (392mg, 65%). NMR: 7.0 (t, 1H), 7.26 (dd, 1H), 7.34 (t, 2H), 7.73 (d, 2H), 8.46 (d, 1H), 8.69 (dd, 1H), 8.76 (s, 1H), 9.67 (s, 1H), 10.38 (d, 1H).

Example 12**2-(2-Fluoroanilino)-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine**

3-(3-Dimethylaminoprop-2-en-1-yl)imidazo[1,2b]pyridazine (Method 17, 1.0g, 4.63mmol), 2-fluorophenylguanidine hydrogen carbonate (Method 25; 1.1g, 5.09mmol) and sodium methoxide (550mg, 10.2mmol) were suspended in DMA (42ml) and the mixture stirred and heated at 140°C for 5.5 hours. The mixture was allowed to cool to ambient temperature and then acetic acid (0.33ml) was added and the volatiles removed by evaporation. The residue was triturated with water and the product collected by filtration, washed with water and dried to give title compound (406mg, 29 %) as a brown solid. NMR: 7.13 (m, 1H), 7.2 (m, 2H), 7.4 (dd, 1H), 7.93 (m, 2H), 8.8 (d, 1H), 8.43 (s, 1H), 8.58 (d, 1H), 8.77 (d, 1H), 9.05 (s, 1H); m/z: 307 [MH]⁺.

Example 13**2-Anilino-4-(2-methylimidazo[1,2b]pyridazin-3-yl)pyrimidine**

3-(3-Dimethylaminoprop-2-en-1-yl)-2-methylimidazo[1,2b]pyridazine (Method 18; 1.5g, 6.52mmol), phenylguanidine hydrogen carbonate (1.41g, 7.17mmol) and sodium methoxide (704mg, 13.04mmol) were suspended in dimethylacetamide (62ml) and the mixture stirred and heated at 160°C for 2.5 hours. The mixture was allowed to cool and more phenylguanidine hydrogen carbonate (642mg, 3.25mmol) and sodium methoxide (352mg, 6.52mmol) were added and the reaction mixture heated a further 2 hours at 160°C. The mixture was allowed to cool to ambient temperature and acetic acid (1.16ml) was added and the volatiles removed by evaporation. The triturated with water and the precipitated solid collected by filtration, washed with water, dried and recrystallized from methanol to give the

title compound (785mg, 40%) as a pale yellow solid. NMR: 2.77 (s, 3H), 6.95 (t, 1H), 7.27 (t, 2H), 7.35 (dd, 1H), 7.76 (d, 2H), 7.83 (d, 1H), 8.15 (d, 1H), 8.58 (d, 1H), 8.65 (d, 1H), 9.5 (s, 1H); m/z: 303 [MH]⁺.

5 Examples 14-19

The following compounds were prepared by methods analogous to that described in Example 13.

Ex	Compound	NMR	M/z
14 ¹	2-Anilino-4-(2-ethylimidazo[1,2b]pyridazin-3-yl)pyrimidine	1.27 (t, 3H), 3.27 (q, 2H), 6.93 (t, 1H), 7.27 (m, 2H), 7.36 (dd, 1H), 7.73 (m, 3H), 8.17 (d, 1H), 8.57 (d, 1H), 8.65 (d, 1H), 9.48 (s, 1H)	317 [MH] ⁺
15 ²	2-Anilino-4-(2-dimethylaminoimidazo[1,2b]pyridazin-3-yl)pyrimidine	2.98 (s, 6H), 6.93 (t, 1H), 7.25 (m, 3H), 7.57 (d, 1H), 7.9 (m, 3H), 8.45 (d, 1H), 8.52 (d, 1H), 9.55 (s, 1H)	332 [MH] ⁺
16	2-Anilino-4-(5-methoxyimidazo[1,2b]pyridazin-3-yl)pyrimidine	4.1 (s, 3H), 6.95 (t, 1H), 7.08 (d, 1H), 7.32 (t, 2H), 7.82 (d, 2H), 8.0 (d, 2H), 8.18 (d, 1H), 8.40 (s, 1H), 8.60 (d, 1H), 9.60 (s, 1H)	319 [MH] ⁺
17	2-Anilino-4-(5-(2,2,2-trifluoroethoxy)imidazo[1,2b]pyridazin-3-yl)pyrimidine	5.18 (q, 2H), 6.95 (t, 1H), 7.22 (d, 1H), 7.32 (t, 2H), 7.82 (d, 2H), 7.92 (d, 1H), 8.26 (d, 1H), 8.42 (s, 1H), 8.61 (d, 1H), 9.6 (s, 1H)	387 [MH] ⁺
18	2-Anilino-4-(6,7-dimethylimidazo[1,2b]pyridazin-3-yl)pyrimidine	2.39 (s, 3H), 2.58 (s, 3H), 6.95 (t, 1H), 7.34 (t, 2H), 7.88 (d, 2H), 7.92 (d, 1H), 8.45 (s, 1H), 8.58 (m, 2H), 9.58 (s, 1H)	317 [MH] ⁺
19	2-Anilino-4-(6-methylimidazo[1,2b]pyridazin-3-yl)pyrimidine	2.45 (s, 3H), 6.96 (t, 1H), 7.32 (t, 2H), 7.87 (d, 2H), 7.92 (d, 1H), 8.08 (s, 1H), 8.48 (s, 1H), 8.58 (d, 1H), 9.48 (s, 1H)	303 [MH] ⁺

¹ Purified by chromatography eluting with dichloromethane / methanol (98.5 : 1.5) NMR;;
m/z:.

² Purified by chromatography eluting with dichloromethane / methanol (98 : 2)

5 **Example 20**

2-(2-Fluoro-4-[N-(2-methoxyethyl)sulphamoyl]anilino)-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine

Chlorosulphonic acid (109µl, 1.64mmol) was added to a stirred solution of 2-(2-fluoroanilino)-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine (Example 12; 130mg, 0.41mmol) in
10 thionyl chloride (2.6ml) cooled to 0°C. The reaction was stirred at 0°C for 5 minutes, then heated at 90°C for 80 minutes, allowed to cool and the volatiles removed by evaporation. 2-Methoxyethylamine (0.75ml, 8.58mmol) in methanol (2.0ml) was added to the residue and the mixture stirred at ambient temperature for 20 hours. The volatiles were removed by
evaporation, water added to the residue and the mixture stirred for 30 minutes. The resulting
15 precipitate was collected by filtration, washed with water and dried to give the title compound (217mg, 86 %) as a white solid. NMR: 2.94 (t, 2H), 3.17 (s, 3H), 3.28 (t, 2H), 7.43 (dd, 1H), 7.65 (dd, 3H), 8.06 (d, 1H), 8.32 (d, 1H), 8.41 (t, 1H), 8.55 (s, 1H), 8.66 (d, 1H), 8.78 (d, 1H), 9.45 (s, 1H); m/z: 444 [MH]⁺.

20 **Example 21**

2-(2-Fluoro-4-sulphamoylanilino)-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine

2-(2-Fluoroanilino)-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine (Example 12; 175mg, 0.572mmol) was treated by the method in Example 20 and reacted with a 2.0M solution of ammonia in methanol (4.3ml) to give the title compound (112mg, 51%) as a white solid.
25 NMR: 7.33 (s, 2H), 7.45 (dd, 1H), 7.67 (m, 2H), 8.07 (d, 1H), 8.33 (m, 2H), 8.53 (s, 1H), 8.66 (d, 1H), 8.8 (d, 1H), 9.45 (s, 1H); m/z: 386 [MH]⁺.

Examples 22-26

The following compounds were prepared by a method analogous to that described in Example 21.

Ex	Compound	NMR	M/z
22 ¹	2-(4-Sulphamoylanilino)-4-(2-methylimidazo[1,2b]pyridazin-3-yl)pyrimidine	2.82 (s, 3H), 7.13 (s, 2H), 7.37 (dd, 1H), 7.73 (d, 2H), 7.9 (d, 1H), 7.97 (d, 2H), 8.16 (d, 1H), 8.65 (m, 2H), 9.97 (s, 1H)	380 [M-H] ⁻
23 ²	2-(4-Sulphamoylanilino)-4-(2-ethylimidazo[1,2b]pyridazin-3-yl)pyrimidine	1.30 (t, 3H), 3.27 (q, 2H), 7.13 (s, 2H), 7.38 (dd, 1H), 7.73 (d, 2H), 7.87 (d, 1H), 7.95 (d, 2H), 8.2 (d, 1H), 8.65 (m, 2H), 9.97 (s, 1H)	396 [MH] ⁺
24 ³	2-(4-Sulphamoylanilino)-4-(5-methoxyimidazo[1,2b]pyridazin-3-yl)pyrimidine	4.19 (s, 3H), 7.09 (d, 1H), 7.13 (s, 2H), 7.78 (d, 2H), 8.0 (d, 2H), 8.06 (d, 1H), 8.2 (d, 1H), 8.43 (s, 1H), 8.68 (d, 1H)	398 [MH] ⁺
25 ⁴	2-(4-Sulphamoylanilino)-4-(5-(2,2,2-trifluoroethoxy)imidazo[1,2b]pyridazin-3-yl)pyrimidine	5.2 (q, 2H), 7.15 (s, 2H), 7.24 (d, 1H), 7.78 (d, 2H), 7.98 (d, 2H), 8.02 (d, 1H), 8.3 (d, 1H), 8.5 (s, 1H), 8.7 (d, 1H), 10.04 (s, 1H)	466 [MH] ⁺
26	2-(4-Sulphamoylanilino)-4-(6,7-dimethylimidazo[1,2b]pyridazin-3-yl)pyrimidine	2.38 (s, 3H), 2.47 (s, 3H), 7.12 (s, 2H), 7.78 (d, 2H), 7.97 (d, 1H), 8.0 (d, 2H), 8.5 (s, 1H), 8.6 (s, 1H), 8.65 (d, 1H), 10.02 (s, 1H)	396 [MH] ⁺

5 ¹ Purified by chromatography on a amine Isolute column eluting with methanol / dichloromethane (1 : 1)

² Purified by trituration with dichloromethane / methanol

³ Purified by chromatography on silica gel chromatography eluting with dichloromethane / methanol (99 : 1) increasing polarity to (80 : 20)

10 ⁴ Purified by chromatography a 10g amino Isolute column eluting with dichloromethane / methanol (1 : 1)

Examples 27-38

The following compounds were prepared by a method analogous to that described in Example 20.

Ex	Compound	NMR	M/z
27	2-{4-[N-(2-Methoxyethyl)sulphamoyl]anilino}-4-(2-methylimidazo[1,2b]pyridazin-3-yl)pyrimidine	(DMSO _d ₆ / CD ₃ CO ₂ D): 2.8 (s, 3H), 2.87 (t, 2H), 3.15 (s, 3H), 3.3 (t, 2H), 7.36 (dd, 1H), 7.7 (d, 2H), 7.92 (d, 1H), 7.98 (d, 2H), 8.15 (d, 1H), 8.64 (m, 2H), 9.97 (s, 1H)	438 [M-H] ⁻
28	2-{4-[N-(3-Dimethylaminopropyl)sulphamoyl]anilino}-4-(2-methylimidazo[1,2b]pyridazin-3-yl)pyrimidine	1.47 (m, 2H), 2.03 (s, 6H), 2.13 (t, 2H), 2.73 (t, 2H), 2.8 (s, 3H), 7.36 (m, 2H), 7.67 (d, 2H), 7.92 (d, 1H), 8.0 (d, 2H), 8.17 (d, 1H), 8.67 (m, 2H), 10.0 (s, 1H)	465 [M-H] ⁻
29	2-[4-(N-Methylsulphamoyl)anilino]-4-(2-methylimidazo[1,2b]pyridazin-3-yl)pyrimidine	2.4 (d, 3H), 2.8 (s, 3H), 7.17 (q, 1H), 7.37 (dd, 1H), 7.67 (d, 2H), 7.9 (d, 1H), 8.0 (d, 2H), 8.16 (d, 1H), 8.65 (m, 2H), 10.0 (s, 1H)	394 [M-H] ⁻
30	2-{4-[N-(2-Methoxyethyl)sulphamoyl]anilino}-4-(2-ethylimidazo[1,2b]pyridazin-3-yl)pyrimidine	1.32 (t, 3H), 2.87 (t, 2H), 3.15 (s, 3H), 3.27 (m, 4H), 7.38 (dd, 1H), 7.45 (s, 1H), 7.7 (d, 2H), 7.88 (d, 1H), 7.98 (d, 2H), 8.2 (d, 1H), 8.65 (d, 2H), 10.0 (s, 1H)	453 [MH] ⁺
31	2-[4-(N-Methylsulphamoyl)anilino]-4-(2-ethylimidazo[1,2b]pyridazin-3-yl)pyrimidine	1.33 (t, 3H), 2.38 (d, 3H), 3.25 (q, 2H), 7.2 (q, 1H), 7.38 (dd, 1H), 7.68 (d, 2H), 7.9 (d, 1H), 7.98 (d, 2H), 8.2 (d, 1H), 8.63 (m, 2H), 9.97 (s, 1H)	410 [MH] ⁺
32 1	2-{4-[N-(2-Methoxyethyl)sulphamoyl]anilino}-4-(2-dimethylaminoimidazo[1,2b]pyridazin-3-yl)pyrimidine	2.85 (t, 2H), 2.97 (s, 6H), 3.17 (m, 5H), 7.3 (dd, 1H), 7.45 (t, 1H), 7.68 (m, 3H), 7.97 (dd, 1H), 8.1 (d, 2H), 8.49 (dd, 1H), 8.58 (d, 1H), 10.08 (s, 1H)	469 [MH] ⁺

33 ²	2-{4-[N-(2-Dimethylaminoethyl)sulphamoyl]anilino}-4-(5-methoxyimidazo[1,2b]pyridazin-3-yl)pyrimidine	2.23 (s, 6H), 2.82 (t, 2H), 3.0 (m, 2H), 4.08 (s, 3H), 7.08 (d, 1H), 7.58 (br s, 1H), 7.98 (d, 2H), 8.06 (d, 2H), 8.1 (d, 1H), 8.44 (s, 1H), 8.68 (d, 1H), 10.12 (s, 1H)	469 [MH] ⁺
34	2-[4-(N-Methylsulphamoyl)anilino]-4-(5-methoxyimidazo[1,2b]pyridazin-3-yl)pyrimidine	2.41 (d, 3H), 4.08 (s, 3H), 7.08 (d, 1H), 7.2 (m, 1H), 7.73 (d, 2H), 8.04 (d, 2H), 8.08 (d, 1H), 8.19 (d, 1H), 8.45 (s, 1H), 8.64 (d, 1H), 10.07 (br s, 1H)	412 [MH] ⁺
35 ³	2-{4-[N-(2-Methoxyethyl)sulphamoyl]anilino}-4-(5-methoxyimidazo[1,2b]pyridazin-3-yl)pyrimidine	2.9 (t, 2H), 3.18 (s, 3H), 3.28 (m, 2H) 4.1 (s, 3H), 7.12 (d, 1H), 7.43 (t, 1H), 7.76 (d, 2H), 8.04 (d, 2H), 8.1 (d, 1H), 8.2 (d, 1H), 8.43 (s, 1H), 8.68 (d, 1H), 10.07 (s, 1H)	456 [MH] ⁺
36	2-{4-[N-(3-Methoxypropyl)sulphamoyl]anilino}-4-(5-methoxyimidazo[1,2b]pyridazin-3-yl)pyrimidine	1.6 (quin, 2H), 2.78 (q, 2H), 3.17 (s, 3H), 3.25 (t, 2H), 4.08 (s, 3H), 7.08 (d, 1H), 7.32 (t, 1H), 7.75 (d, 2H), 8.02 (d, 2H), 8.08 (d, 1H), 8.19 (d, 1H), 8.46 (s, 1H), 8.66 (d, 1H), 10.07 (s, 1H)	470 [MH] ⁺
37 ⁴	2-{4-[N-(2-Methoxyethyl)sulphamoyl]anilino}-4-(6,7-dimethylimidazo[1,2b]pyridazin-3-yl)pyrimidine	2.37 (s, 3H), 2.56 (s, 3H), 2.9 (t, 2H), 3.16 (s, 3H), 3.31 (t, 2H), 7.35 (t, 1H), 7.75 (d, 2H), 8.01 (d, 1H), 8.05 (d, 2H), 8.52 (s, 1H), 8.48 (s, 1H), 8.64 (d, 1H), 9.9 (s, 1H)	454 [MH] ⁺
38 ⁵	2-{4-[N-(2-Methoxyethyl)sulphamoyl]anilino}-4-(6-methylimidazo[1,2b]pyridazin-3-yl)pyrimidine	2.49 (s, 3H), 2.92 (q, 2H), 3.19 (s, 3H), 3.34 (q, 2H), 7.47 (t, 1H), 7.78 (d, 2H), 8.04 (d, 1H), 8.09 (d, 2H), 8.14 (s, 1H), 8.6 (s, 1H), 8.69 (d, 1H), 8.74 (d, 1H), 10.1 (s, 1H)	440 [MH] ⁺
39 ⁶	2-{4-[N-(2-Ethoxyethyl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	1.04 (t, 3H), 2.90 (q, 2H), 3.35 (m, 4H), 7.44 (m, 2H), 7.76 (d, 2H), 8.08 (m, 3H), 8.36 (d, 1H), 8.66 (s, 1H), 8.70 (d, 1H), 8.80 (d, 1H), 10.11 (s, 1H)	440 [MH] ⁺

¹ Purified by reverse phase chromatography on a Hypersil BDS C18 21 X 100mm column eluting with acetonitrile / water / trifluoroacetic acid (10 : 90 : 0.1) decreasing in polarity to (90 : 10 : 0.1)

² Triturated with methanol / ether / dichloromethane

5 ³ Purified by chromatography on silica gel eluting with dichloromethane / methanol (95 : 5) and then with dichloromethane / 2M methanolic ammonia solution (95 : 5)

⁴ Triturated with methanol

⁵ Purified by chromatography on silica gel eluting with dichloromethane / methanol (97 : 3)

⁶ Purified by further trituration with methanol

10

Example 40

2-{4-[N-(1-Methylcycloprop-1-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine

Chlorosulphonic acid (0.22ml, 3.3mmol) was added to a solution of 2-anilino-4-
15 (imidazo[1,2b]pyridazin-3-yl)pyrimidine (Example 10; 300mg, 1.04mmol) in thionyl chloride (6ml) cooled at 5°C. The mixture was stirred at 5°C for 30 minutes, then at ambient temperature for 1.5 hours and then heated at reflux for 1.5 hours. 1-Amino-1-methylcyclopropane (Method 48; 2ml) in isopropanol (10ml) and dimethylethylamine (3ml) was added and the mixture stirred at ambient temperature for 18 hours. The solvent was
20 evaporated and the residue purified by chromatography on silica gel eluting with dichloromethane / isohexane (1 : 1) increasing in polarity to dichloromethane / methanol (95 : 5). The purified product was triturated with ether / methanol to give the title compound (89mg, 21%). NMR: 0.16-0.19 (m, 2H), 0.40-0.44 (m, 2H), 0.92 (s, 3H), 7.25 (dd, 1H), 7.55-7.60 (m, 3H), 7.86-7.90 (m, 3H), 8.15 (d, 1H), 8.48 (s, 1H), 8.53 (d, 1H), 8.62 (d, 1H), 9.95 (s,
25 1H); m/z: 422 [MH]⁺.

Examples 41-57

The following compounds were prepared by a method analogous to that described in Example 40.

Ex	Compound	NMR	M/z
41	2-{4-[<i>N</i> -(3-Methoxypropyl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	1.58 (quin, 2H), 2.78 (q, 2H), 3.14 (s, 3H), 3.22-3.25 (m, 2H), 7.32 (t, 1H), 7.44 (dd, 1H), 7.75 (d, 2H), 8.05 (s, 1H), 8.16 (d, 2H), 8.34 (s, 1H), 8.63 (s, 1H), 8.68 (d, 1H), 8.8 (d, 1H), 10.12 (s, 1H)	440 [MH] ⁺
42	2-[4-(<i>N</i> -Propylsulphamoyl)anilino]-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	0.79 (t, 3H), 1.38 (q, 2H), 2.69 (q, 2H), 7.32 (t, 2H), 7.45 (dd, 1H), 7.75 (d, 2H), 8.04 (s, 1H), 8.09 (d, 2H), 8.34 (d, 1H), 8.63 (s, 2H), 8.68 (d, 1H), 8.80 (d, 1H), 10.11 (s, 1H)	410 [MH] ⁺
43	2-[4-(<i>N</i> -Ethylsulphamoyl)anilino]-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	0.96 (t, 3H), 2.78 (q, 2H), 7.30 (t, 1H), 7.43 (dd, 1H), 7.75 (d, 2H), 8.05 (s, 1H), 8.08 (d, 2H), 8.35 (d, 1H), 8.64 (s, 1H), 8.69 (d, 1H), 8.80 (d, 1H), 10.11 (s, 1H)	396 [MH] ⁺
44	2-[4-(<i>N</i> -Allylsulphamoyl)anilino]-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	3.40 (t, 2H), 5.01 (d, 1H), 5.15 (d, 1H), 5.61-5.70 (m, 1H), 7.45 (dd, 1H), 7.56 (t, 1H), 7.78 (d, 2H), 8.04 (s, 1H), 8.07 (d, 2H), 8.35 (d, 1H), 8.64 (s, 1H), 8.70 (d, 2H), 8.80 (d, 1H), 10.12 (s, 1H)	408 [MH] ⁺
45	2-[4-(<i>N</i> -Propargylsulphamoyl)anilino]-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	3.64 (s, 2H), 7.44 (dd, 1H), 7.78 (d, 2H), 8.08 (m, 3H), 8.34 (d, 1H), 8.70 (d, 1H), 8.80 (d, 1H), 10.13 (s, 1H)	406 [MH] ⁺
46	2-{4-[<i>N</i> -(2-(1-Pyrazolyl)ethyl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	3.11 (q, 2H), 4.15 (t, 2H), 6.20 (s, 1H), 7.39 (s, 1H), 7.44 (dd, 1H), 7.59 (t, 1H), 7.65 (s, 1H), 7.74 (s, 2H), 8.08 (s, 1H), 8.09 (d, 2H), 8.32 (d, 1H), 8.64 (s, 1H), 8.69 (d, 1H), 8.80 (d, 1H), 10.13 (s, 1H)	462 [MH] ⁺

47	2-{4-[<i>N</i> -(1,3-Dihydroxyprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	3.01-3.06 (m, 1H), 3.25-3.35 (m, 4H), 4.55 (t, 2H), 7.18 (d, 1H), 7.49 (dd, 1H), 7.80 (dd, 2H), 8.04 (s, 1H), 8.09 (dd, 2H), 8.35 (d, 1H), 8.68 (s, 1H), 8.70 (d, 1H), 8.80 (d, 1H), 10.15 (s, 1H)	442 [MH] ⁺
48	2-{4-[<i>N</i> -(1-Hydroxyprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	0.88 (d, 3H), 3.05-3.12 (m, 2H), 3.20-3.38 (m, 1H), 4.60 (s, 1H), 7.24 (d, 1H), 7.44 (dd, 1H), 7.77 (d, 2H), 8.04-8.08 (m, 3H), 8.35 (d, 1H), 8.64 (s, 1H), 8.68 (d, 1H), 8.80 (d, 1H), 10.11 (s, 1H)	
49	2-{4-[<i>N</i> -(1-Methoxyprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	0.90 (d, 3H), 3.05-3.15 (m, 1H), 3.15 (s, 3H), 3.23-3.25 (m, 2H), 7.40 (d, 1H), 7.45 (dd, 1H), 7.78 (d, 2H), 8.04-8.09 (m, 3H), 8.35 (d, 1H), 8.64 (s, 1H), 8.70 (d, 1H), 8.80 (d, 1H), 10.15 (s, 1H)	
50	2-{4-[<i>N</i> -(2-Methylprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	1.09 (s, 9H), 7.22 (s, 1H), 7.44 (dd, 1H), 7.78 (d, 2H), 8.03 (s, 1H), 8.08 (d, 2H), 8.34 (d, 1H), 8.64 (s, 1H), 8.70 (d, 1H), 8.80 (d, 1H), 10.09 (s, 1H)	422 [MH] ⁺
51	2-{4-[<i>N</i> -(2-Dimethylamino-ethyl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	2.08 (s, 6H), 2.28 (t, 2H), 2.82 (t, 2H), 7.24 (s, 1H), 7.42 (dd, 1H), 7.79 (d, 2H), 8.06 (s, 1H), 8.10 (d, 2H), 8.35 (d, 1H), 8.64 (s, 1H), 8.70 (d, 1H), 8.80 (d, 1H), 10.11 (s, 1H)	439 [MH] ⁺
52	2-[4-(<i>N</i> -cyclobutylsulphamoyl)anilino]-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	1.42-1.50 (m, 2H), 1.75 (q, 2H), 1.88-1.95 (m, 2H), 3.56-3.64 (m, 1H), 7.45 (dd, 1H), 7.66-7.78 (m, 3H), 8.02-8.08 (m, 3H), 8.35 (d, 1H), 8.64 (s, 1H), 8.70 (d, 1H), 8.80 (d, 1H), 10.11 (s, 1H)	422 [MH] ⁺

53	2-{4-[<i>N</i> -(1,3-Dihydroxy-2-methylprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	0.98 (s, 3H), 3.25-3.31 (m, 4H), 4.52 (t, 2H), 6.72 (s, 1H), 7.45 (dd, 1H), 7.80 (d, 2H), 8.02-8.10 (m, 3H), 8.35 (d, 1H), 8.64 (s, 1H), 8.70 (d, 1H), 8.80 (d, 1H), 10.10 (s, 1H)	456 [MH] ⁺
54	2-[4-(<i>N</i> -cyclopropylmethylsulphamoyl)anilino]-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	0.01-0.08 (m, 2H), 0.26-0.30 (m, 2H), 0.70-0.79 (m, 1H), 2.59 (t, 2H), 7.40 (dd, 1H), 7.72 (d, 2H), 8.0 (s, 1H), 8.02 (d, 2H), 8.30 (d, 1H), 8.60 (s, 2H), 8.64 (d, 1H), 8.78 (d, 1H), 10.05 (s, 1H)	422 [MH] ⁺
55	2-[4-(<i>N</i> -cyclopropylsulphamoyl)anilino]-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	0.01-0.08 (m, 2H), 0.10-0.15 (m, 2H), 1.72-1.78 (m, 1H), 7.10 (dd, 1H), 7.30 (s, 1H), 7.40 (d, 1H), 7.70-7.78 (m, 3H), 8.0 (d, 1H), 8.33 (s, 1H), 8.35 (d, 1H), 8.46 (d, 1H), 9.8 (s, 1H)	408 [MH] ⁺
56	2-{4-[<i>N</i> -(2-Methoxyethyl)- <i>N</i> -methylsulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	2.71 (s, 3H), 3.10 (t, 2H), 3.22 (s, 3H), 3.42 (t, 2H), 7.42 (dd, 2H), 7.75 (d, 2H), 8.08 (d, 1H), 8.12 (d, 2H), 8.32 (d, 1H), 8.68 (s, 1H), 8.70 (d, 1H), 8.80 (d, 1H), 10.18 (s, 1H)	440 [MH] ⁺
57	2-{4-[<i>N</i> -(1-Hydroxy-2-methylprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	1.2 (s, 6H), 3.20 (d, 2H), 4.75 (t, 1H), 7.09 (s, 1H), 7.49 (dd, 1H), 7.82 (d, 2H), 8.05 (d, 2H), 8.09 (d, 1H), 8.38 (d, 1H), 8.68 (s, 1H), 8.70 (d, 1H), 8.85 (d, 1H), 10.15 (s, 1H)	440 [MH] ⁺

Example 58

2-{4-[*N*-(1-Methoxy-2-methylprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine

5

Sodium methoxide (96g 1.8mmol) was added to 2-{4-[*N*-(2,2-dimethylaziridyl)sulphonyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine (Method 46; 150mg, 0.36mmol) in methanol (5ml) and the mixture heated at 50°C for 1.5 hours. The solvent was removed by evaporation, the residue dissolved in ethyl acetate / methanol, washed with water, dried (MgSO₄) and the solvent removed by evaporation. The crude product was purified by reverse phase chromatography to give the title compound (19mg, 12%). NMR: 1.12 (s, 6H),

10

1.88 (m, 2H), 3.15 (s, 3H), 7.42 (d, 1H), 7.77 (dd, 2H), 8.0-8.05 (m, 3H), 8.27 (d, 1H), 6.25 (s, 1H), 8.64 (d, 1H), 8.74 (d, 1H); m/z: 454 [MH]⁺.

Example 59

5 2-{4-[N-(2-Methoxy-2-methylprop-1-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine

The second fraction from chromatographic purification described in Example 58 gave the title compound (10mg, 7%). NMR: 1.09 (s, 6H), 2.65 (d, 2H), 3.05 (s, 3H), 7.25 (t, 1H), 7.46 (dd, 1H), 7.78 (d, 2H), 8.05-8.10 (m, 3H), 8.30 (d, 1H), 8.64 (s, 1H), 8.68 (d, 1H), 8.80 (d, 10 1H), 10.1 (s, 1H); m/z: 454 [MH]⁺.

Example 60

2-{4-[N-(1-(Pyrrolidin-1-yl)-2-methylprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine

15 2-{4-[N-(2,2-Dimethylaziridyl)sulphonyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine (Method 46; 150mg, 0.35mmol) in pyrrolidine (4ml) was stirred and heated at 50°C for 2 hours and then at, ambient temperature for 65 hours. The volatiles were evaporated and the residue dissolved in ethyl acetate / methanol, washed with water and dried (Na₂SO₄). The solvent was removed by evaporation and the residue triturated with ether to give the title 20 compound (96mg, 55%). NMR: 1.05 (s, 6H), 1.58-1.63 (m, 4H), 2.43 (s, 2H), 2.53-2.58 (m, 4H), 6.96 (s, 1H), 7.44 (dd, 1H), 7.79 (d, 2H), 8.05 (d, 2H), 8.07 (d, 1H), 8.36 (d, 1H), 8.64 (s, 1H), 8.69 (d, 1H), 8.80 (d, 1H), 10.08 (s, 1H); m/z: 493 [MH]⁺.

Example 61

25 2-{4-[N-(1-Methylthio-2-methylprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine

A mixture of 2-{4-[N-(2,2-dimethylaziridyl)sulphonyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine (Method 46; 150mg, 0.36mmol) and sodium methanethiolate (125mg, 1.8mmol) in DMF (4ml) was heated at 80°C for 18 hours. The 30 solvent was removed by evaporation, the residue dissolved in ethyl acetate / methanol, washed with water, dried (MgSO₄) the solvent evaporated. The residue was triturated with ether to give the title compound (90mg, 54%). NMR: 1.15 (s, 6H), 2.10 (s, 3H), 2.65 (s, 2H), 7.40 (s,

1H), 7.50 (dd, 1H), 7.82 (d, 2H), 8.08-8.1 (m, 3H), 8.39 (d, 1H), 8.69 (s, 1H), 8.73 (d, 1H), 8.80 (d, 1H), 10.15 (s, 1H); m/z: 470 [MH]⁺.

Example 62

5 **2-{4-[N-(1-Morpholino-2-methylprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine**

2-{4-[N-(2,2-Dimethylaziridyl)sulphonyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine (120mg, 0.29mmol) in morpholine (5ml) was heated reflux 3 hours. The solvent was evaporated, the residue was dissolved in ethyl acetate / methanol, washed with water,
10 dried (MgSO₄) and the solvent evaporated. The residue was triturated with ether to give the title compound (50mg, 35%). NMR: 1.05 (s, 6H), 2.29 (s, 2H), 2.43-2.50 (m, 4H), 3.48-3.52 (m, 4H), 7.04 (s, 1H), 7.43 (dd, 1H), 7.78 (d, 2H), 8.02-8.08 (m, 3H), 8.35 (d, 1H), 8.64 (s, 1H), 8.71 (d, 1H), 8.80 (d, 1H), 10.08 (s, 1H).

15 **Example 63**

2-{4-[N-(1,3-Dimethoxyprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine

Chlorosulphonic acid (0.23ml, 3.3mmol) was added to a solution of 2-anilino-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine (Example 10; 300mg, 1.04mmol) in thionyl chloride
20 (6ml) at 5°C. The mixture was stirred at 5°C for 30 minutes, then at ambient temperature for 1 hour and finally heated at reflux for 1.5 hour. The mixture was allowed to cool to ambient temperature and a solution of 1,3-dimethoxy-2-aminopropane (Method 57; 14mmol) in ethanol (20ml) and dimethylethylamine (1ml) were added to the residue, and the mixture stirred at ambient temperature for 18 hours. The volatiles were removed by evaporation, the
25 residue triturated with water and the solid product collected by filtration and then dried under vacuum at 60°C to give the title compound (322mg, 66%). NMR: 3.10 (s, 6H), 3.21 (d, 4H), 7.42-7.58 (m, 2H), 7.78 (d, 2H), 8.02-8.08 (m, 3H), 8.35 (d, 1H), 8.65-8.70 (m, 2H), 8.80 (d, 1H), 10.09 (s, 1H); m/z: 470 [MH]⁺.

Examples 64-66

The following compounds were prepared by a method analogous to that described in Example 63.

Ex	Compound	NMR	M/z
64 ¹	2-{4-[N-(1-Ethoxyprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	0.90 (d, 3H), 1.05 (t, 3H), 3.08-3.12 (m, 1H), 3.22-3.30 (m, 4H), 7.39 (d, 2H), 7.45 (dd, 1H), 7.79 (d, 2H), 8.03-8.09 (m, 3H), 3.35 (d, 1H), 8.64 (s, 1H), 8.70 (d, 1H), 8.80 (d, 1H)	454 [MH] ⁺
65 ²	2-{4-[N-(1-Propoxyprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	0.79 (t, 3H), 0.94 (d, 3H), 1.40 (q, 2H), 3.08-3.12 (m, 1H), 3.18-3.20 (m, 2H), 7.38 (d, 1H), 7.43 (dd, 1H), 7.78 (d, 2H), 8.03-8.09 (m, 3H), 8.36 (d, 1H), 8.62 (s, 1H), 8.68 (d, 1H), 8.81 (d, 1H), 10.11 (s, 1H)	468 [MH] ⁺
66 ³	2-{4-[N-(1,3-Diethoxyprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine	0.99 (t, 6H), 3.23-3.31 (m, 9H), 7.42-7.51 (m, 2H), 7.78 (d, 2H), 8.02-8.09 (m, 3H), 8.34 (d, 1H), 8.62 (s, 1H), 8.70 (d, 1H), 8.80 (d, 1H), 10.08 (s, 1H)	498 [MH] ⁺

- ¹ 1-Ethoxy-2-aminopropane (Method 54) as starting material, purified by chromatography
5 eluting with dichloromethane / isohexane (1 : 1) increasing in polarity to dichloromethane / methanol (97 : 3)
² 1-Propoxy-2-aminopropane (Method 55) as starting material, purified by chromatography on silica gel eluting with dichloromethane / methanol (100 : 0) increasing in polarity to (95 : 5).
³ 1,3-Diethoxy-2-aminopropane (Method 56) as starting material, purified by chromatography
10 on silica gel eluting with dichloromethane / isohexane (1 : 1) increasing in polarity to dichloromethane / methanol (98.5 : 1.5)

Preparation of Starting Materials

- The starting materials for the Examples above are either commercially available or are
15 readily prepared by standard methods from known materials. For example the following reactions are illustrations but not limitations of the preparation of some of the starting materials used in the above reactions.

Method 1**3-(1-Hydroxyethyl)imidazo[1,2a]pyrazine**

Anhydrous neutral alumina (activity 1) (28.5g) was added to a stirred solution of 2-aminopyrazine (2.85g, 30mmol) in dry dichloromethane (100ml) under nitrogen. A solution of 2,3-epoxybutanal (5.58g, 30mmol) in dichloromethane (10ml) was added and the reaction mixture stirred for 18 hours. The reaction mixture was filtered, and the alumina cake washed with dichloromethane (100ml), and methanol/dichloromethane (1:1, 100ml). The filtrates were combined, the volatiles were removed by evaporation (keeping the bath temperature below 40°C). The resulting crude yellow oil was triturated with dichloromethane and the solid collected by filtration to give the title compound (586mg, 12%) as a pale orange solid. NMR: 1.6 (d, 3H), 5.18 (quin, 1H), 5.5 (d, 1H), 7.7 (s, 1H), 7.88 (d, 1H), 8.5 (d, 1H), 9.0 (s, 1H).

Method 2**3-Acetylimidazo[1,2a]pyrazine**

3-(1-Hydroxyethyl)imidazo[1,2a]pyrazine (Method 1; 3.87g, 23.7mmol) was dissolved in acetone (30ml) with manganese dioxide (21.1g, 237mmol) and stirred at ambient temperature for 72 hours. The solution was filtered through diatomaceous earth, washed with acetone (4 x 50ml) and then with methanol (3 x 50ml). The solvent was removed from the filtrate by evaporation. The residue was purified by column chromatography eluting with methanol/dichloromethane (0:100 increasing in polarity to 10:90) to give the title compound (349mg, 10%) as a pure crystalline white solid. NMR: 2.60 (s, 1H), 8.22 (d, 1H), 8.75 (s, 1H), 9.35 (m, 2H).

Method 3**3-(3-Dimethylaminoprop-2-en-1-yl)imidazo[1,2a]pyrazine**

A mixture of 3-acetylimidazo[1,2a]pyrazine (Method 2; 340mg, 2.11mmol) and DMFDMA (20ml) were heated under reflux, under nitrogen for 18 hours. The mixture was allowed to cool and the solvent removed by evaporation. The residue was triturated with ether, collected by filtration and dried to give the title compound (406mg, 89%) as a dark red solid. NMR: 2.91 (br s, 3H) 3.15 (br s, 3H), 5.4 (d, 1H), 7.75 (d, 1H), 8.18 (d, 1H), 8.6 (s, 1H), 9.19 (s, 1H), 9.55 (dd, 1H); m/z: 218 [MH]⁺.

Method 4**2-[4-Sulphonoylanilino]-4-(imidazo[1,2a]pyrazin-3-yl)pyrimidine**

2-Anilino-4-(imidazo[1,2a]pyrazin-3-yl)pyrimidine (Example 9; 350mg, 1.22mmol) was added in portions to concentrated sulphuric acid (6ml) stirred and cooled with an ice-bath to maintain temperature <25°C. The reaction mixture was allowed to warm to ambient temperature and stirred for 1 hour. The mixture was then carefully poured into ice-water (150ml) and the ice allowed to melt. The sodium hydroxide pellets and then 1M aqueous sodium hydroxide solution was carefully added to adjusted the solution from pH0.5 to pH2. The mixture was stirred at ambient temperature for 18 hours and the resulting precipitate collected by filtration, washed with water (2 x 15ml) and dried. The crude product was triturated with dichloromethane then recollected by filtration and then was re-triturated with dichloromethane/methanol (1:1) recollected by filtration and dried to give the title compound (191mg, 52%). NMR: 7.54 (d, 1H), 7.60 (d, 2H), 7.64 (d, 2H), 8.15 (d, 1H), 8.54 (d, 1H), 8.81 (s, 1H), 9.95 (br s, 1H); m/z: 367 [M-H]⁺.

Method 5**3-(1-Hydroxyethyl)imidazo[1,2a]pyrimidine**

2-Aminopyrimidine (2.85g, 30mmol) was treated by the method described in Method 1 to give crude product which was purified by column chromatography on neutral alumina (activity 3) eluting with dichloromethane/methanol (100:0 increasing in polarity to 95:5) to give the title compound (662mg, 14%). NMR: 1.59 (d, 3H), 5.12 (quin, 1H), 5.39 (d, 1H), 7.15 (dd, 1H), 7.60 (s, 1H), 8.53 (dd, 1H), 8.96 (dd, 1H); m/z: 164 [MH]⁺.

Method 6**3-Acetylimidazo[1,2a]pyrimidine**

3-(1-Hydroxyethyl)imidazo[1,2a]pyrimidine (Method 5; 630mg, 3.87mmol) was treated with manganese dioxide (3.44g, 38.7mmol) as described in Method 2 but without chromatographic purification to give the title compound (379mg, 60%) as a pale yellow solid. NMR: 2.56 (s, 3H), 7.29 (t, 1H), 8.76 (s, 1H), 8.80 (m, 1H), 9.78 (dd, 1H).

Method 7**3-(3-Dimethylaminoprop-2-en-1-yl)imidazo[1,2a]pyrimidine**

3-Acetylimidazo[1,2a]pyrimidine (Method 6; 370mg, 2.29mmol) was treated as described in Method 3 to give the title compound (479mg, 96%). NMR: 2.94 (br s, 1H), 3.15 (br s, 3H), 5.83 (d, 1H), 7.22 (dd, 1H), 7.70 (s, 1H), 8.60 (s, 1H), 8.68 (dd, 1H), 8.96 (dd, 1H); m/z: 217 [MH]⁺.

Method 8**2-[4-Sulphonoylanilino]-4-(imidazo[1,2a]pyrimidin-3-yl)pyrimidine**

2-Anilino-4-(imidazo[1,2a]pyrimidin-3-yl)pyrimidine (70% isomeric mixture with 2-anilino-4-(imidazo[1,2a]pyrimidin-2-yl)pyrimidine) (Example 11; 392mg, 1.36mmol) was treated as described in Method 4 to give the title compound as the major component (70%) in an isomeric mixture with 2-[4-sulphonoylanilino]-4-(imidazo[1,2a]pyrimidin-2-yl)pyrimidine (363mg, 73%). NMR: 7.47 (d, 1H), 7.45-7.7 (m, 5H), 8.57 (d, 1H), 8.93 (dd, 1H), 9.02 (s, 1H), 9.82 (s, 1H), 10.4 (d, 1H).

Method 9**5-Chloroimidazo[1,2b]pyridazine**

A mixture of 3-amino-6-chloropyridazine (5.0g, 38.6mmol) and chloroacetaldehyde (5.9ml of a 7.9M solution in water, 46.3mM) in 1-butanol (37ml) was heated at 120°C with for 20 hours. The reaction was allowed to cool to ambient temperature and the crystallised solid collected by filtration, washed with a little 1-butanol and then ether. The solid was dissolved in water (50ml) and the solution adjusted to pH10 by carefully addition of 40% aqueous sodium hydroxide solution. The resulting suspension was extracted with ethyl acetate. The organic extracts were combined, washed with saturated brine, dried (Na₂SO₄) and the volatiles were removed by evaporation. The residue was triturated with isohexane, collected by filtration, washed with isohexane and dried to give the title compound (4.35g, 74%) as a pale brown solid. NMR: 7.33 (d, 1H), 7.83 (s, 1H), 8.20 (d, 1H), 8.32 (s, 1H); m/z: 154 [MH]⁺.

Method 10Imidazo[1,2b]pyridazine

A mixture of 5-chloroimidazo[1,2b]pyridazine (Method 9; 5.82g, 37.9mmol), triethylamine (5.28ml, 37.9mmol) and 5% palladium on charcoal (300mg) in ethyl acetate (233ml) was stirred vigorously under an atmosphere of hydrogen until the uptake of hydrogen ceased. The reaction mixture was filtered through diatomaceous earth. The filter pad was washed with ethyl acetate and volatiles removed from the filtrate by evaporation. The residue was triturated with ice cold pentane, collected by filtration, washed sparingly with cold pentane and dried to give the title compound (3.65g, 81%) as a white solid. NMR (CDCl₃): 7.03 (dd, 1H), 7.78 (s, 1H), 7.95 (m, 2H), 8.30 (d, 1H); m/z: 120 [MH]⁺.

Method 113-Bromoimidazo[1,2b]pyridazine

N-Bromosuccinimide (2.7g, 15.1mmol) was added to a solution of imidazo[1,2b]pyridazine (Method 10; 1.8g, 15.1mmol) in chloroform (15ml) and the mixture was stirred and heated at reflux for 20 minutes. The reaction mixture was allowed to cool to ambient temperature and saturated aqueous sodium carbonate solution (15ml) was added and the mixture stirred vigorously. More chloroform and water was added to dissolve the precipitated solids. The organic layer was separated, dried (Na₂SO₄) and the solvent removed by evaporation. The residue was recrystallized from ethanol (50ml), collected by filtration, washed with cold ethanol and dried to give the title compound (2.61g, 88%) as an off white solid. NMR (CDCl₃): 7.10 (dd, 1H), 7.80 (s, 1H), 7.95 (d, 1H), 8.47 (d, 1H); m/z: 198 [MH]⁺.

Methods 12-13

The following compounds were prepared by a method analogous to that described in Method 11, except that they were reacted at ambient temperature for 4 hours.

Meth	Compound	NMR	M/z	SM
12	3-Bromo-5-methoxyimidazo[1,2b]pyridazine	3.99 (s, 3H), 6.94 (d, 1H), 7.71 (s, 1H), 8.02 (d, 1H)	230 [MH] ⁺	Meth 41
13 ¹	3-Bromo-5-(2,2,2-trifluoroethoxy)imidazo[1,2b]pyridazine	5.08 (q, 2H), 7.12 (d, 1H), 7.8 (s, 1H), 8.16 (d, 1H)	298 [MH] ⁺	Meth 42

¹ Re-crystallised from ethyl acetate / isohexane

Method 14**3-Acetylimidazo[1,2b]pyridazine**

Isopropyl magnesium bromide (12.5ml of a 1M solution in THF, 12.5mmol) was added dropwise over 5 minutes to a stirred solution of 3-bromoimidazo[1,2b]pyridazine (Method 11; 1.98g, 10mmol) in dry THF (100ml) under nitrogen at -40°C. The reaction mixture was stirred for 105 minutes at -40°C and then dry *N*-methoxy-*N*-methylacetamide (1.65ml, 16mmol) was added. The mixture was stirred a further 30 minutes at -40°C and then allowed to warm to ambient temperature and stirred for a further 90 minutes. Acetic Acid (1.26ml) was added and the volatiles then removed by evaporation. The residue was partitioned between dichloromethane (120ml) and water (30ml). The layers were separated and the aqueous layer was extracted with dichloromethane. The organic layers were combined, dried (Na₂SO₄) and volatiles removed by evaporation. The residue was triturated with ether and isohexane, collected by filtration, washed with isohexane and dried to give the title compound (1.25g, 78%) as a brown solid. NMR (CDCl₃): 2.75 (s, 3H), 7.27 (dd, 1H), 8.08 (d, 1H), 8.45 (s, 1H), 8.62 (d, 1H); m/z: 162 [MH]⁺.

Methods 15-16

The following compounds were prepared by a method analogous to that described in Method 14.

Meth	Compound	NMR	M/z	SM
15 ¹	3-Acetyl-5-methoxyimidazo[1,2b]pyridazine	2.64 (s, 3H), 4.0 (s, 3H), 7.15 (d, 1H), 8.17 (d, 1H), 8.3 (s, 1H)	192 [MH] ⁺	Meth 12
16 ²	3-Acetyl-5-(2,2,2-trifluoroethoxy)imidazo[1,2b]pyridazine	2.65 (s, 3H), 5.10 (q, 2H), 7.30 (d, 1H), 8.27 (d, 1H), 8.37 (s, 1H)		Meth 13

¹ Re-crystallised from ethyl acetate / isohexane

² 1:1 mix of product and 5-(2,2,2-trifluoroethoxy)imidazo[1,2b]pyridazine

Method 17**3-(3-Dimethylaminoprop-2-en-1-oyl)imidazo[1,2b]pyridazine**

3-Acetylimidazo[1,2b]pyridazine (Method 14; 1.25g, 7.76mmol) was suspended in DMFDMA (32ml) and the mixture heated under nitrogen at 100°C for 42 hours. The excess dimethylacetal was removed by evaporation and the residue triturated with ether and isohexane. The product was collected by filtration, washed with isohexane and dried to give the title compound (1.58g, 94%) as a brown solid. NMR (CDCl₃) : 3.00 (s, 3H), 3.17 (s, 3H), 6.20 (d, 1H), 7.13 (dd, 1H), 7.92 (d, 1H), 8.03 (d, 1H), 8.38 (s, 1H), 8.50 (d, 1H); m/z: 217 [MH]⁺.

10

Methods 18-24

The following compounds were prepared by a method analogous to that described in Method 17.

Meth	Compound	NMR	M/z	SM
18	3-(3-Dimethylaminoprop-2-en-1-oyl)-2-methylimidazo[1,2b]pyridazine	2.58 (s, 3H), 2.87 (s, 3H), 2.9 (s, 3H), 6.27 (d, 1H), 7.27 (dd, 1H), 7.73 (d, 1H), 8.05 (d, 1H), 8.56 (d, 1H)	231 [MH] ⁺	Meth 34
19	3-(3-Dimethylaminoprop-2-en-1-oyl)-2-ethylimidazo[1,2b]pyridazine	1.23 (t, 3H), 2.87 (s, 3H), 3.0 (q, 2H), 3.13 (s, 3H), 6.23 (d, 1H), 7.25 (dd, 1H), 7.70 (d, 1H), 8.07 (d, 1H), 8.56 (d, 1H)	245 [MH] ⁺	Meth 35
20	3-(3-Dimethylaminoprop-2-en-1-oyl)-5-methoxyimidazo[1,2b]pyridazine	2.9 (br s, 3H), 3.15 (br s, 3H), 4.02 (s, 3H), 6.43 (d, 1H), 7.0 (d, 1H), 7.8 (d, 1H), 8.03 (s, 1H), 8.05 (d, 1H)	247 [MH] ⁺	Meth 15
21	3-(3-Dimethylaminoprop-2-en-1-oyl)-5-(2,2,2-trifluoroethoxy)imidazo[1,2b]pyridazine	2.94 (br s, 3H), 3.18 (br s, 3H), 5.16 (q, 2H), 6.3 (d, 1H), 7.18 (d, 1H), 7.8 (d, 1H), 8.14 (s, 1H), 8.2 (d, 1H)	315 [MH] ⁺	Meth 16

22	3-(3-Dimethylaminoprop-2-en-1-oyl)-6,7-dimethylimidazo[1,2b]pyridazine	2.35 (s, 3H), 2.50 (s, 3H), 2.95 (br s, 3H), 3.15 (br s, 3H), 6.25 (d, 1H), 7.75 (d, 1H), 8.20 (s, 1H), 8.50 (s, 1H)	245 [MH] ⁺	Meth 37
23	3-(3-Dimethylaminoprop-2-en-1-oyl)-6-methylimidazo[1,2b]pyridazine	2.4 (s, 3H), 2.90 (br s, 3H), 3.15 (br s, 3H), 6.2 (d, 1H), 7.74 (d, 1H), 7.97 (s, 1H), 8.19 (s, 1H), 8.56 (s, 1H)	231 [MH] ⁺	Meth 38
24 ¹	3-(3-Dimethylaminoprop-2-en-1-oyl)-2-dimethylaminoimidazo[1,2b]pyridazine	2.96 (s, 12H), 5.9 (d, 1H), 7.13 (dd, 1H), 7.53 (d, 1H), 7.8 (d, 1H), 8.33 (d, 1H)	260 [MH] ⁺	Meth 36

¹ By a similar method, but heating at 100°C for 120 hours and purifying the product by chromatography eluting with dichloromethane / methanol (95 : 5)

5 Method 25

2-Fluorophenylguanidine bicarbonate

Concentrated hydrochloric acid (6ml) in water (4.8ml) was added to a mixture of 2-fluoroaniline (7.94g, 71.2mmol) and cyanamide (6.98g, 166mmol) and the mixture heated at 115°C for 2.5 hours. The reaction mixture was allowed to cool to ambient temperature and the solution was adjusted to pH 13 by careful addition of 40% aqueous sodium hydroxide solution. The aqueous solution was extracted with ethyl acetate and the combined organic extracts were dried (Na₂SO₄) and the volatiles removed by evaporation. The crude product was dissolved in water (40 ml) and carbon dioxide gas bubbled through the solution until the pH of the suspension remained constant (approximately pH 9). The precipitated solid was collected by filtration, washed sparingly with water and dried to give the title compound (11.95g, 78%) as a white solid. NMR: 6.83 (m, 2H), 7.0 (m, 2H); m/z: 154 [MH]⁺.

Method 26

N,N-Dimethyl-N'-(6-chloropyridazin-3-yl)acetamide

3-Amino-6-chloropyridazine (1.29g, 10mmol) was suspended in dry toluene (20 ml) and dimethylacetamide dimethyl acetal (1.46ml, 10mmol) was added. The mixture was stirred and heated at reflux for 3 hours and then left to stand at ambient temperature for 18 hours.

Insolubles were removed from the reaction mixture by filtration and the filter washed with ethyl acetate. The filtrate was then evaporated and the residue triturated with isohexane, the solid product collected by filtration and washed with isohexane to give the title compound (1.7g, 85%) as a yellow solid. NMR: (CDCl₃) 2.13 (s, 3H), 3.1 (s, 6H), 6.92 (d, 1H), 7.25 (d, 1H); m/z: 199 [MH]⁺.

Methods 27-28

The following compounds were prepared by a method analogous to that described in Method 26.

Meth	Compound	NMR	M/z	SM
27 ¹	<i>N,N</i> -Dimethyl- <i>N'</i> -(6-chloro-4,5-dimethylpyridazin-3-yl)acetamide	2.22 (s, 3H), 2.24 (s, 3H), 3.02 (s, 3H), 3.1 (s, 3H), 8.38 (s, 1H)	213 [MH] ⁺	Com av
28 ²	<i>N,N</i> -Dimethyl- <i>N'</i> -(6-chloro-5-methylpyridazin-3-yl)acetamide	2.24 (s, 3H), 3.0 (s, 3H), 3.12 (s, 3H), 7.1 (s, 1H), 8.45 (s, 1H)	199 [MH] ⁺	Meth 43

¹ Heating at 120°C for 7 hours

² Purified by column chromatography eluting with isohexane / ethyl acetate (70 : 30 increasing in polarity to 90 : 10):-

Method 29

15 3-Acetyl-5-chloro-2-methylimidazo[1,2b]pyridazine

Chloroacetone (2.9ml, 36.2mmol) was added to a suspension of powdered sodium bromide (3.73g, 36.2mmol) in ethanol (75ml) and the mixture stirred at ambient temperature for 2 hours. *N,N*-dimethyl-*N'*-(6-chloropyridazin-3-yl)acetamide (Method 26; 6.0g, 30.2mmol) was added and the suspension stirred and heated at reflux for 5.5 hours. The reaction mixture was allowed to cool to ambient temperature and the volatiles removed by evaporation. Water (60ml) was added to the residue and the resulting precipitate collected by filtration, washed with water and dried to give the title compound (4.93g, 79%) as a pale yellow solid. NMR: 2.6 (s, 3H), 2.7 (s, 3H), 7.57 (d, 1H), 8.23 (d, 1H); m/z: 209 [MH]⁺.

Methods 30-33

The following compounds were prepared by a method analogous to that described in Method 29.

Meth	Compound	NMR	M/z	SM
30	3-Acetyl-5-chloro-2-ethylimidazo[1,2b]pyridazine	1.23 (t, 3H), 2.7 (s, 3H), 3.02 (q, 2H), 7.58 (d, 1H), 8.25 (d, 1H)	224 [MH] ⁺	Meth 39
31 ¹	3-Acetyl-5-chloro-2-dimethylaminoimidazo[1,2b]pyridazine	(CDCl ₃): 2.78 (s, 3H), 3.13 (s, 6H), 7.13 (d, 1H), 7.63 (d, 1H)	239 [MH] ⁺	Meth 40
32 ²	3-Acetyl-5-chloro-6,7-dimethylimidazo[1,2b]pyridazine	2.4 (s, 3H), 2.62 (s, 3H), 2.65 (s, 3H), 8.50 (s, 1H)	224 [MH] ⁺	Meth 27
33 ³	3-Acetyl-5-chloro-6-methylimidazo[1,2b]pyridazine	2.45 (s, 3H), 2.6 (s, 3H), 8.32 (s, 1H), 8.52 (s, 1H)	210 [MH] ⁺	Meth 28

¹ Using acetonitrile as the reaction solvent and heating at 65°C for 24 hours:-

5 ² Using DMF as the reaction solvent and heating at 70°C for 12 hours followed by purification of the product by passing through an SCX ion exchange column eluting with 2N methanolic ammonia / dichloromethane (50 : 50) and recrystallisation from ethyl acetate / methanol

³ Using DMF as the reaction solvent and sodium iodide in place of sodium bromide and heating at 70°C for 12 hours

10

Method 34**3-Acetyl-2-methylimidazo[1,2b]pyridazine**

A mixture of 3-acetyl-5-chloro-2-methylimidazo[1,2b]pyridazine (Method 29; 4.8g, 22.9mmol), triethylamine (3.21ml, 22.9mmol) and 5% palladium on charcoal (1.6g) in ethyl acetate (180ml) was stirred vigorously under an atmosphere of hydrogen until the uptake of hydrogen ceased. The reaction mixture was filtered through diatomaceous earth. The filter pad was washed with ethyl acetate and volatiles removed from the filtrate by evaporation. The residue was triturated with isohexane, collected by filtration, washed sparingly with isohexane and dried to give the title compound (3.7g, 92%) as a white solid. NMR: 2.6 (s, 3H), 2.73 (s, 3H), 7.43 (dd, 1H), 8.15 (d, 1H), 8.67 (d, 1H); m/z: 176 [MH]⁺.

15

20

Methods 35-38

The following compounds were prepared by a method analogous to that described in Method 34.

Meth	Compound	NMR	M/z	SM
35	3-Acetyl-2-ethyl imidazo[1,2b]pyridazine	1.23 (t, 3H), 2.73 (s, 3H), 3.02 (q, 2H), 7.43 (dd, 1H), 8.18 (d, 1H), 8.68 (d, 1H)	190 [MH] ⁺	Meth 30
36	3-Acetyl-2-dimethylamino imidazo[1,2b]pyridazine	2.67 (s, 3H), 3.0 (s, 6H), 7.35 (dd, 1H), 7.93 (d, 1H), 8.5 (d, 1H)	205 [MH] ⁺	Meth 31
37	3-Acetyl-6,7-dimethyl imidazo[1,2b]pyridazine	2.39 (s, 3H), 2.55 (s, 3H), 2.65 (s, 3H), 8.45 (s, 1H), 8.78 (s, 1H)	190 [MH] ⁺	Meth 32
38 ¹	3-Acetyl-6-methyl imidazo[1,2b]pyridazine	2.44 (s, 3H), 2.62 (s, 3H), 8.07 (q, 1H), 8.44 (s, 1H), 8.64 (d, 1H)	176 [MH] ⁺	Meth 33

¹ Using ethanol as the reaction solvent

5

Method 39*N,N*-Dimethyl-*N'*-(6-chloropyridazin-3-yl)propionamidine

A solution of phosphorous oxytrichloride (9.32ml, 100mmol) in dry toluene (30ml) was added dropwise over 15 minutes to a stirred solution of dimethyl propionamide (10.88ml, 100mmol) in dry toluene (180ml) and the mixture was then stirred for 24 hours at ambient temperature. 3-Amino-6-chloropyridazine (12.9g, 100mmol) was added as a solid and the mixture stirred and heated at 85°C for 18 hours. The reaction mixture was allowed to cool and settle and the toluene layer decanted off. The residue was washed with more dry toluene and this decanted off. The residue was dissolved in water and extracted with dichloromethane and the aqueous layer was adjusted to pH 12 by careful addition of 40% aqueous sodium hydroxide solution. Dichloromethane was added and the mixture was filtered to remove the insoluble impurity. The organic layer was separated and the aqueous layer extracted with dichloromethane. The organic extracts were combined, dried (Na₂SO₄) and the volatiles removed by evaporation. The crude product was purified by column chromatography on silica gel eluting with dichloromethane / methanol (96.5 : 3.5) to give the title compound (4.4g, 21%) as a brown oil. NMR (CDCl₃): 1.18 (t, 3H), 2.53 (q, 2H), 3.1 (s, 6H), 6.93 (d, 1H), 7.27 (d, 1H); m/z: 213 [MH]⁺.

20

Method 40

The following compound was prepared by a method analogous to that described in Method 39.

Meth	Compound	NMR	M/z
40	<i>N,N,N',N'</i> -Tetramethyl- <i>N''</i> -(6-chloropyridazin-3-yl)guanidine	(CDCl ₃) 3.0 (s, 12H), 7.5 (d, 1H), 7.67 (d, 1H)	228 [MH] ⁺

5 **Method 41**5-Methoxyimidazo[1,2b]pyridazine

Sodium methoxide solution (25% wt solution in methanol, 56.5g, 261.4mmol) was added to a solution of 5-chloroimidazo[1,2b]pyridazine (Method 9; 10.0g, 65.6mmol) in anhydrous methanol (100ml) at ambient temperature and the reaction mixture stirred for 18 hours. The volatiles were removed by evaporation and the yellow, oily residue was dissolved in dichloromethane (100ml). The solution was washed with water (5 x 100ml) until aqueous wash became neutral. The organic solution was dried (MgSO₄) and the solvent removed to give the title compound (8.87 g, 91%) as a pale yellow solid. NMR: 3.92 (s, 3H), 6.82 (d, 1H), 7.59 (s, 1H), 7.99 (d, 1H), 8.03 (s, 1H); m/z: 150 [MH]⁺.

15

Method 425-(2,2,2-Trifluoroethoxy)imidazo[1,2b]pyridazine

Sodium hydride (432mg, 10.8mmol) was added in portions to a solution of 2,2,2-trifluoroethanol (10.8mmol) and 5-chloroimidazo[1,2b]pyridazine (1.5g, 9.8mmol), in anhydrous DMF (15ml) at ambient temperature and the mixture stirred for 18 hours. The volatiles were removed by evaporation and the resulting solid was dissolved in dichloromethane (20ml). The solution was washed with water (2 x 15ml), the organic layer dried (MgSO₄) and the solvent removed by evaporation to give the title compound (2.05g, 92%) as a pale yellow solid. NMR: 5.0 (q, 2H), 7.0 (d, 1H), 7.64 (s, 1H), 8.1 (m, 2H); m/z: 219 [MH]⁺.

25

Method 43**3-Amino-6-chloro-5-methylpyrazine**

A mixture of 3,6-dichloro-5-methylpyrazine (5g, 30.7mmol) in ethanolic ammonia solution (50ml) was heated for 8 hours in a high-pressure cell at 135°C. The solvent was then removed by evaporation and the residue dissolved in chloroform (40ml). The organic solution was washed with water (2 x 25ml) dried (MgSO₄) and the volatiles removed by evaporation. The crude solid was semi-purified by column chromatography on silica gel eluting with dichloromethane / methanol (95 : 5) to the title compound (1.3g, 31%) as a 10:17 mixture with its regioisomer. NMR: 2.06 (s, 3H), 2.18 (s, 3H), 6.2 (br s, 4H), 6.74 (s, 1H), 7.26 (s, 1H); m/z: 143 [MH]⁺.

Method 44**1-(2-Aminoethyl)pyrazole**

Sodium hydroxide (22.96g, 0.57mol) was added to a solution of pyrazole (10.88g, 0.16mol) in dry acetonitrile (80ml) and the mixture stirred at ambient temperature for 30 minutes. Tetrabutylammonium hydrogen sulphate (2.18g, 6.4mmol) and 2-chloroethylamine hydrochloride (19.78g, 0.172 mole) were added and the mixture heated at reflux for 24 hours. The mixture was allowed to cool and the insolubles were removed by filtration. The volatiles were removed from the filtrate by evaporation and 49% hydrogen bromide (30ml) followed by ethanol (100ml) added to the residue. The mixture was heated to reflux and then cooled in ice. The resulting precipitate was collected by filtration, washed with cold ethanol to give the title compound. NMR: 3.20-3.24 (m, 2H), 4.38 (t, 2H), 6.30 (s, 1H), 7.50 (s, 1H), 7.79 (s, 1H), 7.98 (s, 2H).

Method 45**2-{4-[N-(1-(4-Toluenesulphonyloxy)-2-methylprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine**

4-Toluenesulphonyl chloride (4.44g, 0.23mol) was added to a solution of 2-{4-[N-(1-hydroxy-2-methylprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine (Example 57; 3.41g, 0.008mol) in pyridine (50ml) and the mixture stirred at ambient temperature for 24 hours. The mixture was then diluted with water and sonicated. The aqueous phase was decanted and the residue dissolved in ethyl acetate / methanol. The solution was washed with water, dried (MgSO₄), the solvent removed by evaporation and the

residue was triturated with ether to give the title compound (1.92g, 42%). NMR: 1.0 (s, 6H), 2.38 (s, 3H), 3.79 (s, 2H), 7.40-7.45 (m, 3H), 7.59 (s, 1H), 7.65-7.75 (m, 4H), 8.05 (d, 2H), 8.09 (d, 1H), 8.35 (d, 1H), 8.66 (s, 1H), 8.70 (d, 1H), 8.80 (d, 1H), 10.13 (s, 1H).

5 Method 46

2-{4-[N-(2,2-dimethylaziridyl)sulphonyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine

Potassium carbonate (57mg, 0.4mmol) and then acetone (5ml) was added to 2-{4-[N-(1-(4-toluenesulphonyloxy)-2-methylprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine (Method 45; 220mg, 0.38mmol) and the mixture stirred and heated at reflux for 2 hours. The mixture was allowed to cool, the insolubles were removed by filtration, the filter washed with acetone, and the solvent removed by evaporation to give the title compound (123mg, 77%) as an orange solid. NMR: 1.42 (s, 6H), 2.43 (s, 2H), 7.43 (dd, 1H), 7.84 (d, 2H), 8.08-8.12 (m, 3H), 8.35 (d, 1H), 8.68 (d, 1H), 8.70 (d, 1H), 8.8 (d, 1H), 10.2 (s, 1H).

15 Method 47

1-(1-Methylcyclopropane)carboxamide

Oxalyl chloride (8.24ml, 0.095mol) and then DMF (few drops) were added to a solution of 1-(1-methylcyclopropane)carboxylic acid (9.42g, 0.094mol) in dichloromethane (150ml) cooled at 5°C and the mixture stirred at 5°C for 30 minutes and then for 3 hours at ambient temperature. The solvent and excess oxalyl chloride were removed by evaporation, the residue dissolved in dichloromethane and added to a solution of ammonia (excess) in methanol cooled at 5°C. The mixture was allowed to warm to ambient temperature and the volatiles removed by evaporation to give the title compound. NMR: 0.29 (q, 2H), 0.71 (q, 2H), 1.02 (s, 3H), 6.62 (s, 1H), 6.85 (s, 1H).

Method 48

1-Amino-1-methylcyclopropane

Bromine (2.87ml, 0.056mol) was added to a solution of sodium hydroxide (13.5g, 0.338mol) in water (100ml) at 0-5°C. A slurry of 1-(1-methylcyclopropane)carboxamide (5.70g 0.056mol) in water (50ml) was then added and reaction mixture stirred at 5°C for 2 hours, then left to stand at ambient temperature for 24 hours. The mixture was then heated at

80°C for 2.5 hours, allowed to cool and mixture distilled to give the title compound (bp 75-80°C). NMR: 0.2 (q, 2H), 0.14 (q, 2H), 0.96 (s, 3H), 1.42 (s, 2H).

Method 49

5 1,3-Dimethoxy-2-methanesulphonyloxypropane

Triethylamine (5ml, 0.036mol) followed by slow addition of methanesulphonyl chloride (2.72ml, 0.035mol) was added to a solution of 1,3-dimethoxy-2-hydroxypropane (3.84g, 0.032mol) in dichloromethane (70ml) cooled at 5°C. The mixture was then stirred at ambient temperature for 24 hours. The mixture was then absorbed onto silica gel and purified
10 by chromatography eluting with dichloromethane / isohexane (1 : 1) to give the title compound (3.74g, 59%). NMR: 3.15 (s, 3H), 3.28 (s, 6H), 3.52 (d, 4H), 4.78 (q, 1H).

Methods 50-52

The following compounds were prepared by a method analogous to that described in
15 Method 49.

Meth	Compound	NMR
50	1-Ethoxy-2-methanesulphonyloxypropane	1.10 (t, 3H), 1.28 (d, 3H), 3.14 (s, 3H), 3.42-3.48 (m, 2H), 3.62-3.68 (m, 2H), 4.78 (q, 1H)
51	1-Propoxy-2-methanesulphonyloxypropane	0.86 (t, 3H), 1.28 (d, 3H), 1.51 (q, 2H), 3.33-3.40 (m, 2H), 3.44 (d, 2H), 3.69 (d, 3H), 4.78 (q, 1H)
52	1,3-Diethoxy-2-methanesulphonyloxypropane	1.12 (t, 6H), 3.16 (s, 3H), 3.41-3.45 (m, 4H), 3.55 (d, 4H), 4.75 (q, 1H)

Method 53

1,3-Dimethoxy-2-azidopropane

1,3-Dimethoxy-2-methanesulphonyloxypropane (Method 49; 3.74g, 19mmol) and
20 sodium azide (2.03g, 31mmol) in DMA (55ml) was heated at 100°C for 8 hours then left to stand at ambient temperature for 24 hours. The mixture was diluted with water, extracted with ethyl acetate, the extracts combined and washed with water, dried (MgSO₄) and the volatiles removed by evaporation to give the title compound (2.0g, 74%) as a clear oil.

Methods 54-56

The following compounds were prepared by a method analogous to that described in Method 53.

Meth	Compound	SM
54	1-Ethoxy-2-azidopropane	Method 50
55	1-Propoxy-2-azidopropane	Method 51
56	1,3-Diethoxy-2-azidopropane	Method 52

5 **Method 57****1,3-Dimethoxy-2-aminopropane**

- 10% Palladium on charcoal (500mg) was added to a solution of 1,3-dimethoxy-2-azidopropane (Method 53; 2g, 0.014mol) in ethanol (40ml) and the mixture stirred under an atmosphere of hydrogen at ambient temperature for 6 hours. The catalyst was removed by
- 10 filtration through diatomaceous earth and the filter pad washed with ethanol to give a solution of the title compound in ethanol (20ml).

Methods 58-60

- The following compounds were prepared by a method analogous to that described in
- 15 Method 57.

Meth	Compound	SM
58	1-Ethoxy-2-aminopropane	Method 54
59	1-Propoxy-2-aminopropane	Method 55
60	1,3-Diethoxy-2-aminopropane	Method 56

Example 67

- The following illustrate representative pharmaceutical dosage forms containing the compound of formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester
- 20 thereof (hereafter compound X), for therapeutic or prophylactic use in humans:-

(a): Tablet I	mg/tablet
Compound X	100
Lactose Ph.Eur	182.75
Croscarmellose sodium	12.0
Maize starch paste (5% w/v paste)	2.25
Magnesium stearate	3.0

(b): Tablet II	mg/tablet
Compound X	50
Lactose Ph.Eur	223.75
Croscarmellose sodium	6.0
Maize starch	15.0
Polyvinylpyrrolidone (5% w/v paste)	2.25
Magnesium stearate	3.0

(c): Tablet III	mg/tablet
Compound X	1.0
Lactose Ph.Eur	93.25
Croscarmellose sodium	4.0
Maize starch paste (5% w/v paste)	0.75
Magnesium stearate	1.0

(d): Capsule	mg/capsule
Compound X	10
Lactose Ph.Eur	488.5
Magnesium stearate	1.5

(e): Injection I	(50 mg/ml)
Compound X	5.0% w/v
1M Sodium hydroxide solution	15.0% v/v
0.1M Hydrochloric acid	(to adjust pH to 7.6)
Polyethylene glycol 400	4.5% w/v
Water for injection	to 100%

(f): Injection II	10 mg/ml
Compound X	1.0% w/v
Sodium phosphate BP	3.6% w/v
0.1M Sodium hydroxide solution	15.0% v/v
Water for injection	to 100%


(g): Injection III	(1mg/ml,buffered to pH6)
Compound X	0.1% w/v
Sodium phosphate BP	2.26% w/v
Citric acid	0.38% w/v
Polyethylene glycol 400	3.5% w/v
Water for injection	to 100%

5 Note

The above formulations may be obtained by conventional procedures well known in the pharmaceutical art. The tablets (a)-(c) may be enteric coated by conventional means, for example to provide a coating of cellulose acetate phthalate.

Claims


1. A compound of formula (I):




(I)

wherein:

Ring A is a group of formula (IA) or (IB):



(IA)



(IB)

- wherein one or more of X^1 , X^2 , X^3 and X^4 are nitrogen and the others are CR^5 wherein the values of R^5 may be the same or different;

- R^1 is halo, nitro, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, amino, carboxy, carbamoyl, mercapto, sulphonamoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkanoylamino, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkylS(O)_a wherein a is 0 to 2, N -(C_{1-6} alkyl)sulphonamoyl or N,N -(C_{1-6} alkyl)₂sulphonamoyl; wherein R^1 may be optionally substituted on carbon by one or more R^6 ;

- R^2 and R^5 are independently of one other selected from hydrogen, halo, nitro, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, amino, carboxy, carbamoyl, mercapto, sulphonamoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkanoylamino, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkylS(O)_a wherein a is 0 to 2, C_{1-6} alkoxycarbonyl, N -(C_{1-6} alkyl)sulphonamoyl, N,N -(C_{1-6} alkyl)₂sulphonamoyl, phenyl, heterocyclic group, phenylthio or (heterocyclic group)thio; wherein any R^2 or R^5 may be optionally substituted on carbon by one or more R^7 ; and wherein if said heterocyclic group contains an $-NH-$ moiety that nitrogen may be optionally substituted by a group selected from R^8 ;

n is 0 to 2, wherein the values of R^1 may be the same or different;

R^3 is halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C_{2-6} alkenyl or C_{2-6} alkynyl;

p is 0-4; wherein the values of R^3 may be the same or different;

5 R^4 is a group A-E-; wherein

A is selected from C_{1-6} alkyl, phenyl, a heterocyclic group, C_{3-8} cycloalkyl, phenyl C_{1-6} alkyl, (heterocyclic group) C_{1-6} alkyl or C_{3-8} cycloalkyl C_{1-6} alkyl; wherein A may be optionally substituted on carbon by one or more R^9 ; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^{10} ;

E is a direct bond or -O-, -C(O)-, -OC(O)-, -C(O)O-, -N(R^a)C(O)-, -C(O)N(R^a)-, -N(R^a)-, -S(O) $_a$ -, -SO $_2$ N(R^a)- or -N(R^a)SO $_2$ -; wherein R^a is hydrogen or C_{1-6} alkyl optionally substituted by one or more R^{11} and a is 0-2;

15 R^9 is independently selected from oxo, halo, nitro, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkylS(O) $_a$ wherein a is 0 to 2, C_{1-6} alkoxycarbonyl, C_{1-6} alkoxycarbonylamino, benzyloxycarbonylamino, N -(C_{1-6} alkyl)sulphamoyl and
20 N,N -(C_{1-6} alkyl) $_2$ sulphamoyl; wherein R^9 may be optionally substituted on carbon by one or more R^{12} ;

q is 0-2; wherein the values of R^4 maybe the same or different; and wherein $p + q \leq 5$;

R^6 , R^7 , R^{11} and R^{12} are independently selected from halo, nitro, cyano, hydroxy, trifluoromethoxy, trifluoromethyl, amino, carboxy, carbamoyl, mercapto, sulphamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxyl, methylamino, ethylamino, dimethylamino, diethylamino, N -methyl- N -ethylamino, acetylamino, N -methylcarbamoyl, N -ethylcarbamoyl, N,N -dimethylcarbamoyl, N,N -diethylcarbamoyl, N -methyl- N -ethylcarbamoyl, methylthio, ethylthio, methylsulphanyl, ethylsulphanyl, mesyl, ethylsulphonyl, methoxycarbonyl, ethoxycarbonyl, N -methylsulphamoyl, N -ethylsulphamoyl, N,N -dimethylsulphamoyl, N,N -diethylsulphamoyl or N -methyl- N -ethylsulphamoyl; and
30

R^8 and R^{10} are independently selected from C_{1-4} alkyl, C_{1-4} alkanoyl, C_{1-4} alkylsulphonyl, C_{1-4} alkoxycarbonyl, carbamoyl, N -(C_{1-4} alkyl)carbamoyl, N,N -(C_{1-4} alkyl)carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl;
or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

5

2. A compound of formula (I) as claimed in claim 1 wherein Ring A is a group of formula (IA) or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

3. A compound of formula (I) as claimed in either of claims 1 or 2 wherein one of X^1 , X^2 , X^3 and X^4 are nitrogen and the others are CR^5 wherein the values of R^5 may be the same or different or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

10

4. A compound of formula (I) as claimed in any one of claims 1-3 wherein n is 0 or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

15

5. A compound of formula (I) as claimed in any one of claims 1-4 wherein R^2 is selected from hydrogen, C_{1-6} alkyl or N,N -(C_{1-6} alkyl)₂amino or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

20

6. A compound of formula (I) as claimed in any one of claims 1-5 wherein R^5 is selected from hydrogen, C_{1-6} alkyl or C_{1-6} alkoxy; wherein R^5 may be optionally substituted on carbon by one or more R^7 ; wherein R^7 is selected from halo or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

25

7. A compound of formula (I) as claimed in any one of claims 1-6 wherein R^3 is sulphamoyl or halo or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

30

8. A compound of formula (I) as claimed in any one of claims 1-7 wherein p is 0 or 1 or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

9. A compound of formula (I) as claimed in any one of claims 1-8 wherein R^4 is a group A-E-; wherein:

A is selected from C_{1-6} alkyl, C_{3-8} cycloalkyl, (heterocyclic group) C_{1-6} alkyl or C_{3-8} cycloalkyl C_{1-6} alkyl; wherein A may be optionally substituted on carbon by one or more R^9 ;

E is $-N(R^a)SO_2-$; wherein R^a is hydrogen or C_{1-6} alkyl;

R^9 is independently selected from hydroxy, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, N,N -(C_{1-6} alkyl) $_2$ amino or C_{1-6} alkylS(O) $_a$ wherein a is 0; or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

10. A compound of formula (I) as claimed in any one of claims 1-9 wherein q is 0 or 1 or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

11. A compound of formula (I) as claimed in any one of claims 1-10 wherein p + q is 0, 1 or 2; wherein the values of R^3 may be the same or different and the values of R^4 may be the same or different; or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

12. A compound of formula (I) as depicted in claim 1 wherein:

Ring A is a group of formula (IA) (as depicted in claim 1)

one of X^1 , X^2 , X^3 and X^4 is nitrogen and the others are CR^5 ; wherein the values of R^5 may be the same or different;

n is 0;

R^2 is selected from hydrogen, methyl, ethyl or N,N -dimethylamino;

R^5 is selected from hydrogen, methyl, methoxy or 2,2,2-trifluoroethoxy;

R^3 is sulphamoyl or fluoro;

p is 0 or 1;

R^4 is N -methylsulphamoyl, N -cyclopropylmethylsulphamoyl, N -ethylsulphamoyl, N -propylsulphamoyl, N -allylsulphamoyl, N -2-propynylsulphamoyl, N -cyclobutylsulphamoyl, N -*t*-butylsulphamoyl, N -cyclopropylsulphamoyl, N -(2-dimethylaminoethyl)sulphamoyl, N -(2-methoxyethyl)sulphamoyl, N -(2-pyrazol-1-ylethyl)sulphamoyl, N -methyl- N -(2-methoxyethyl)sulphamoyl, N -(1-cyclopropylethyl)sulphamoyl,

N-(3-dimethylaminopropyl)sulphamoyl, *N*-(3-methoxypropyl)sulphamoyl,
N-(3-hydroxy-2-hydroxymethylprop-2-yl)sulphamoyl, *N*-(1,3-dihydroxyprop-2-yl)sulphamoyl,
N-(3-morpholino-2-methylprop-2-yl)sulphamoyl, *N*-(1,3-dimethoxyprop-2-yl)sulphamoyl,
N-(1,3-diethoxyprop-2-yl)sulphamoyl, *N*-(1-methoxyprop-2-yl)sulphamoyl,
 5 *N*-(1-ethoxyprop-2-yl)sulphamoyl, *N*-(1-hydroxyprop-2-yl)sulphamoyl,
N-(3-methylthio-2-methylprop-2-yl)sulphamoyl,
N-(3-pyrrolidin-1-yl-2-methylprop-2-yl)sulphamoyl,
N-(3-methoxy-2-methylprop-2-yl)sulphamoyl, *N*-(2-methoxy-2-methylpropyl)sulphamoyl,
N-(1-propoxyprop-2-yl)sulphamoyl or *N*-(3-hydroxy-2-methylprop-2-yl)sulphamoyl;

10 q is 0 or 1;

p + q is 0, 1 or 2; wherein the values of R³ may be the same or different and the values
 of R⁴ may be the same or different;

or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

15 13. A compound of formula (I) as depicted in claim 1 selected from:

2-{4-[N-(2-Methoxyethyl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine;

2-{4-[N-(3-Methoxypropyl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine;

2-[4-(N-Propylsulphamoyl)anilino]-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine;

2-{4-[N-(2-Methoxyethyl)sulphamoyl]anilino}-4-(2-methylimidazo[1,2b]pyridazin-3-

20 yl)pyrimidine;

2-[4-(N-Methylsulphamoyl)anilino]-4-(2-methylimidazo[1,2b]pyridazin-3-yl)pyrimidine;

2-{4-[N-(2-(1-Pyrazolyl)ethyl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-

yl)pyrimidine;

2-{4-[N-(1,3-Diethoxyprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-

25 yl)pyrimidine;

2-{4-[N-(1,3-Dimethoxyprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-

yl)pyrimidine;


2-{4-[N-(1-Ethoxyprop-2-yl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine;

2-{4-[N-(2-Ethoxyethyl)sulphamoyl]anilino}-4-(imidazo[1,2b]pyridazin-3-yl)pyrimidine;

30 or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.


14. A process for preparing a compound of formula (I) as claimed in claim 1 or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof which process (wherein variable groups are, unless otherwise specified, as defined in claim 1) comprises of:

a) reaction of a pyrimidine of formula (II):




(II)

wherein L is a displaceable group; with an amine of formula (III):




(III)

10 b) reacting a pyrimidine of formula (IV):



(IV)


with a compound of the formula (V):



(V)

wherein one of M and Q is a displaceable group X and the other is an metallic reagent Y; or


c) reacting a compounds of formula (VI):



- 72 -

(VI)


with a compound of formula (VII):



(VII)


5 wherein R⁵ is C₁₋₆alkyl and R⁶ is hydrogen or R¹;

d) reacting an amino compound of formula (VIII):



(VIII)

with a compound of formula (IX):



10

wherein E is a displaceable group;

and thereafter if necessary:

i) converting a compound of the formula (I) into another compound of the formula (I);

ii) removing any protecting groups;

15 iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

15. A pharmaceutical composition which comprises a compound of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in any one of claims 1-13 in association with a pharmaceutically-acceptable diluent or carrier.

20

16. A compound of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in any one of claims 1-13 for use as a medicament.

17. The use of a compound of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined in any one of claims 1-13 in the manufacture of a medicament for use in the production of a cell cycle inhibitory (anti-cell-proliferation) effect in a warm-blooded animal such as man.

18. A method of producing a cell cycle inhibitory (anti-cell-proliferation) effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof as claimed in any one of claims 1-13.

19. A pharmaceutical composition which comprises a compound of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in any one of claims 1-13 in association with a pharmaceutically-acceptable diluent or carrier for use in the production of a cell cycle inhibitory (anti-cell-proliferation) effect in a warm-blooded animal such as man.

20. A compound of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in any one of claims 1-13 for use in the production of a cell cycle inhibitory (anti-cell-proliferation) effect in a warm-blooded animal such as man.

INTERNATIONAL SEARCH REPORT

national Application No

PCT/GB 02/00603

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D487/04 A61K31/5025 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 33798 A (DOHERTY ANNETTE MARIAN ; DOBRUSIN ELLEN MYRA (US); WARNER LAMBERT C) 6 August 1998 (1998-08-06) claims 20,27,28; example 17; table 4	1-20
X,P	WO 01 14375 A (BEATTIE JOHN FRANKLIN ; BREAUULT GLORIA ANNE (GB); JEWSBURY PHILLIP) 1 March 2001 (2001-03-01) page 1, line 27 -page 2, line 3; claim 1	1-20

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

17 April 2002

Date of mailing of the international search report

24/04/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Seelmann, I

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 02/00603

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9833798	A	06-08-1998	AU 6648098 A	25-08-1998
			BR 9807305 A	02-05-2000
			EP 0964864 A2	22-12-1999
			HR 980060 A1	30-06-1999
			JP 2001509805 T	24-07-2001
			WO 9833798 A2	06-08-1998
			ZA 9800914 A	09-11-1998
WO 0114375	A	01-03-2001	AU 6583300 A	19-03-2001
			WO 0114375 A1	01-03-2001